

Accepted and scheduled for publication in *The Astrophysical Journal*, for the March 20, 2013, v 766, 1st issue

Stability of the photon indices in Z-source GX 340+0 for spectral states

Elena Seifina¹, Lev Titarchuk² & Filippo Frontera³

ABSTRACT

We show an analysis of the spectral and timing properties of X-ray radiation from Z-source GX 340+0 during its evolutions when the electron temperature of the *Transition Layer* (TL) kT_e monotonically decreases from 21 to 3 keV. We analyze episodes observed with the *BeppoSAX* and *Rossi X-ray Timing Explorer* (*RXTE*). We reveal that the X-ray broad-band energy spectra during all spectral states can be reproduced by a physical model, composed of a soft *Blackbody* component, two Comptonized components (both due to the presence of that TL that up-scatters both seed photons of $T_{s1} \lesssim 1$ keV coming from the disk (first component *Comptb1*), and seed photons of temperature $T_{s2} \lesssim 1.5$ keV coming from the neutron star (second component *Comptb2*) and the iron-line (*Gaussian*) component. Spectral analysis using this model indicates that the photon power-law indices Γ_{com1} and Γ_{com2} of the Comptonized components are almost constant, Γ_{com1} and $\Gamma_{com2} \sim 2$ when kT_e changes from 3 to 21 keV along Z-track. We interpret the detected quasi-stability of the indices of Comptonized components near a value of 2. Furthermore, this index stability now found for the Comptonized spectral components of Z-source GX 340+0 is similar to that was previously established in the *atoll* sources 4U 1728-34, GX 3+1 and early proposed for a number of X-ray neutron stars (NSs). This behavior of NSs both for *atoll* and Z-sources, is essentially different from that observed in black hole binaries where Γ_{com} increases during a spectral evolution from the low state to the high state and ultimately saturates at high mass accretion rate.

¹Moscow State University/Sternberg Astronomical Institute, Universitetsky Prospekt 13, Moscow, 119992, Russia; seif@sai.msu.ru

²Dipartimento di Fisica, Università di Ferrara, Via Saragat 1, I-44122 Ferrara, Italy, email:titarchuk@fe.infn.it; George Mason University Fairfax, VA 22030; Goddard Space Flight Center, NASA, code 663, Greenbelt MD 20770, USA; email:lev@milkyway.gsfc.nasa.gov, USA

³Dipartimento di Fisica, Università di Ferrara, Via Saragat 1, I-44122 Ferrara, Italy, email:frontera@fe.infn.it

Subject headings: accretion, accretion disks—stars: neutron—black hole physics—stars: individual (GX 340+0):radiation mechanisms: nonthermal—physical data and processes

1. Introduction

The study of binaries hosting neutron stars (NSs) allows one to investigate properties of compact objects themselves and to better understand the differences between black hole (BH) and NS systems. Low-mass X-ray binaries (LMXBs) including weakly magnetized neutron stars may be observationally divided into two main sub-classes, so called *Z* and *atoll* sources, based upon correlations between their spectral and timing properties during state evolution in X-ray wavelengths (Hasinger & van der Klis 1989). The similarity and differences between *atoll* and *Z* sources are clear visible on the *color-color diagram* (CCD) or *hardness-intensity diagram* (HID) and determined by their timing properties.

The *Z*-shaped track shown by *Z*-sources in the CCD or HID consist of three branches: the *horizontal*, the *normal* and the *flaring* branches (HB, NB and FB respectively). Note the aforementioned names are merely historical [see e.g. van der Klis (1989)] and were based on the fact that, in the first few sources studied, the horizontal branch appeared as a horizontally oriented limb in the HIDs, the normal branch as a state in which the sources pass most of their time, while in the flaring branch the sources demonstrated intensive flares. In turn, *atoll* sources showed two kinds of branches: a *banana* shaped branch and an *island* state. Note that *atoll* sources usually show their pattern over a larger extent of luminosity and on longer time scales (weeks - month) than that of *Z*-sources (hours - weeks). *Atoll* sources always follow *banana* branch on the same short time scale as *Z*-sources, hours - weeks (Muno et al. 2002).

The occurrence of the different X-ray states of the *Z*-sources and the *atoll* sources is presumably governed by mass accretion rate in their innermost parts of the source. In the *Z*-sources this mass accretion rate increases from HB, through NB, to FB (Hasinger et al. 1990, Vrtilek et al. 1990). On the other hand in the *atoll* sources mass accretion rate increases from the *island* state, through the lower *banana* to the upper *banana* (see e.g. Corbet et al. 1989; Di Salvo et al., 2001). Hasinger & van der Klis (1989) showed that each state has a characteristic fast timing variability, and therefore an extra diagnostic is available to identify a source state, and to distinguish the *Z*-sources from the *atoll* sources. In fact, only a combined diagnostic including fast timing and spectral analysis (such as CCDs) can provide reliable identification of the source (*Z* or *atoll*) type .

The division of the bright LMXBs into *Z* and *atoll* sources was proposed because of differences in both magnetic fields and mass accretion rate (Hasinger & van der Klis 1989). In fact, for *Z*-sources higher magnetic fields were suggested ($10^9 - 10^{10}$ G) than that of *atoll* sources ($10^8 - 10^9$ G) and they probably accrete at near-Eddington rates (e.g. Lamb 1989), while mass accretion rate $\dot{M}$ is well below Eddington rates in *atoll* sources. This conjecture on the magnetic field is quite old-dated and no evidence has ever been given. Furthermore *the discovery of the first transient Z source, turning to an atoll makes the B-field discriminant very unlikely* [see Homan et al. (2007); Lin et al. (2009)]. A study of broad band spectral and timing behavior turns to be a powerful approach to probe the Physics of the brightest LMXBs. This study is also used to investigate the behavior of other X-ray binaries, e.g. black hole binaries (see e.g. van der Klis 1994). An overall comparison of these various types of sources shows that the properties of the rapid fluctuations appear to depend on $\dot{M}$ in the innermost part of the source and a compact object type (BH or NS).

Usually BH and NS sources demonstrate state transitions when the soft X-ray flux monotonically increases (or decreases) which implies monotonic changes of $\dot{M}$ [e.g., Titarchuk & Seifina (2009), Seifina & Titarchuk (2011) and below]. However, the total X-ray flux is not at all times a reliable indicator for $\dot{M}$ (e.g., Mendez et al. 1999; Ford et al. 2000).

A *Z*-type represents a category of seven Galactic LMXBs: Sco X-1, GX 17+2, GX 349+2, Cyg X-2, GX 5-1, GX 340+0 and XTE J1701-462 (with an addition of Cir X-1 which is a *peculiar Z*-source, see Shirey et al., 1999) and the extra-galactic *Z*-source LMC X-2. It has been suggested that there are two types of *Z*-sources, characterized by differences in the shape of their *Z*-patterns and their power spectral characteristics (Hasinger & van der Klis 1989, see also Penninx et al. 1991).

GX 340+0 was discovered as a Galactic X-ray source by Friedman, Byram & Chubb (1967) and later its position was found by Rappaport et. al (1971). This source was preliminary determined as a NS binary by Margon et al. (1971) using an *Aerobee* rocket data. *Ariel* (Ponman, 1982), *EXOSAT* (van Paradijs et al., 1988), HEAO-2 (Schulz et al. 1989) orbital observatories provided unique high quality X-ray data of GX 340+0 to indicate three main tracks with characteristic correlation of hardness ratio, which formed *Z*-pattern in CCD. Despite of no burst has yet been found, the very high values of QPO ($\gtrsim 800$ Hz) and X-ray energy spectrum (van Paradijs et al., 1988) and as well as the similarities with the different *Z*-sources allowed to classify GX 340+0 as a *Z*-source (Hasinger & Van der Klis, 1989).

The spectral characteristics of this source have not been fully studied up to now. Schulz & Wijers (1993), hereafter SW93, carried out a spectral analysis of the source based on *EXOSAT* data in the 2 – 12 keV energy range. They found that the spectrum can be fit by model which a sum of a blackbody of a temperature of 0.8 keV and a thermal Comptonization

component of soft NS seed photons. SW93 also obtained an optical depth of $\tau \sim 5 - 6$ and an electron temperature of the hot corona changed from 4 to 6 keV.

Fender & Hendry (2000) estimated the distance to the source of 11 ± 3 kpc using the radio measurements. Radio emission of GX 340+0 is highly variable, and probably correlates with the X-ray emission on its horizontal branch. But the radio emission anti-correlates when the source is on the other spectral branches (Oosterbroek et al. 1994).

Lavagetto et al. (2004) were the first who showed a broadband (0.1 – 200 keV) spectrum of this source based on the data using a *BeppoSAX* observation on August 9 – 10, 2001 (thereafter Observation *S6*, see Table 1). They decomposed the spectrum of GX 340+0 as a soft blackbody of temperature $kT_{BB} \sim 0.5$ keV, Comptonization component and in addition an excess at photon energies more than 20 keV was fit using an extended power law terminated by a cutoff at high energies. Ueda et al. (2005) studied a spectrum of the source using *Chandra* observations. These *Chandra* data indicated to the presence of an iron line emission. The shape of the line can be fit by a *Gaussian* profile, which center is at 6.6 keV and the line equivalent width is 40 eV. Then, D’Ai et al. (2009) detected a broad asymmetric emission line in the Fe K_α energy band and tried to investigate the line evolution along HB of *Z*-track based on XMM-*Newton* data. However, the line profile does not indicate to any correlated variation although the continuum emission and luminosity underwent significant changes.

Iaria et al. (2006), Iaria06 hereafter, also examined the 0.1 – 200 keV energy spectrum using *BeppoSAX* observations (thereafter Observations *S3* – *S6*, see Table 1) and the Lavagetto model described above. For all observations of different spectral states, they found $N_H \sim 6.2 \times 10^{22} \text{ cm}^{-2}$ and that the spectrum at less than 30 keV could be fit by a sum a blackbody of $kT_{BB} \sim 0.5$ keV and a Comptonization component *Comptt* (Titarchuk, 1994) which seed-photon temperature of $kT_s \sim 0.9$ keV and plasma temperature of $kT_e \sim 3$ keV. In addition, Iaria06 used a power-law component of the photon index $\Gamma \sim 1.5$ and a *Gaussian* line component at 6.75 keV, related to Fe XXV. They also found that a power-law component of a photon index of 2.5 was required at higher energies.

Church et al. (2006) investigated the 3–40 keV spectrum relied on PCA/*RXTE* data (*R3* set in terms of our classification, see Table 2) and applied a modified Lavagetto model, with a *cutoff power law* at the energies above 30 keV. They showed a good agreement this model with *RXTE* observations at a stable value of the best-fit photon index $\Gamma \sim 1.7$ during all spectral states.

Before the stability of the index at different luminosities was emphasized by Farinelli & Titarchuk (2011) (hereafter FT11) for a number of neutron star low mass X-ray binaries (LMXBs).

FT11 gathered X-ray *BeppoSAX* spectra and showed the stability of the spectral index α at approximately 1 ($\Gamma = \alpha + 1$) for many NSs: X 1658-298, GX 354-0, GS 1826-238, 1E 1724-3045, Cyg X-2, Sco X-1, GX 17+2, and GX 349+2 at their different spectral states (or when the best-fit electron temperatures for these sources are drastically different). X-ray spectra for *atoll* sources 4U 1728-34 and GX 3+1 detected with *BeppoSAX* and *RTXE*, respectively, at different luminosities were studied in details by Seifina & Titarchuk (2011), (2012), hereafter ST11, and ST12 respectively.

The ST11 and ST12 also pointed out that the value of Γ only varies slightly around 2, independently of kT_e and luminosity. This stability of Γ can be an universal signature for any kind NSs either the *atoll* or *Z*-sources. It is presumably dictated by common Physics for this type of sources. FT11, ST11 and ST12 demonstrated this *stability* of Γ using a model in which the emergent spectrum influenced by a strong Comptonization component is produced in the Transition Layer (TL) between NS surface and the *Keplerian* accretion disk. In fact, *the stability of the index occurs when Q_{cor} , which is the energy release in TL, is much higher than Q_{disk} which is the photon energy flux, coming from the accretion disk and illuminating TL.*

Namely, ST11 shows (see also FT11) that the TL optical depth τ_0 , its electron temperature kT_e , the ratio Q_{disk}/Q_{cor} are related to each other

$$\frac{kT_e\tau_0(2 + \tau_0)}{m_e c^2} = \frac{0.25}{1 + Q_{disk}/Q_{cor}}, \quad (1)$$

if one assumes that the balance of energy in the transition layer (TL) is established by gravitational energy release and cooling of electrons due to inverse Compton effect and free-free emission [see Zel'dovich & Shakura (1969) and later a similar formulation of this problem in Bisnovatyi et al. (1980)].

The spectral index α of the Comptonization spectrum is determined by the formula

$$\alpha = -\frac{3}{2} + \sqrt{\frac{9}{4} + \frac{\beta}{\Theta}}, \quad (2)$$

where the dimensionless plasma temperature $\Theta \equiv kT_e/m_e c^2$ and β -parameter [see Titarchuk & Lyubarskij (1995)]. If one replaces β by its diffusion approximation

$$\beta_{diff} = \frac{1}{\tau_0(2 + \tau_0)} \quad (3)$$

and applies equation (1), we have a formula for the diffusion spectral index α_{diff}

$$\alpha_{diff} = -\frac{3}{2} + \sqrt{\frac{9}{4} + \frac{1 + Q_{disk}/Q_{cor}}{0.25}}, \quad (4)$$

and the photon index

$$\Gamma_{diff} \approx 1 + \alpha_{diff} = 2 + 0.8 Q_{disk}/Q_{cor} \quad (5)$$

for $Q_{disk}/Q_{cor} < 1$. Consequently $\Gamma_{diff} \approx 2$ as $Q_{disk}/Q_{cor} \ll 1$.

Note, in contrast with the neutron star sources many black hole (BH) candidate binaries show a correlated behavior of the photon index Γ with mass accretion rate $\dot{M}$ (see Shaposhnikov & Titarchuk 2009 and Titarchuk & Seifina 2009, hereafter ST09 and TS09, respectively).

Previous investigations (see Penninx et al. 1991; Kuulkers & van der Klis 1996; Jonker et al. 1998) show that *Z*-sources, in particular GX 340+0, are characterized, in HB or in an upper part of NB, by QPOs with frequencies changing from 20 to 50 Hz. van Paradijs et al. (1988) revealed normal branch oscillations (NBOs) with a frequency of 5.6 Hz in the middle of NB. Jonker et al. (1998) found twin kHz QPOs whose centroid frequencies are correlated with the HBO (HB oscillation) frequency.

Titarchuk & Osherovich (2001) classified the power density spectrum features for each compact object class using the *Transition Layer* model [see Titarchuk et al. (1998)] and the basic properties of the accretion flow around compact object. They applied their analysis for the *RXTE* observations of GX 340+0.

It is interesting that other *Z*-sources differ from GX 340+0 on the timing properties along *Z*-track. In particular, Cyg X-2 and GX 5-1 have well developed HBs with a strong HBO and red low-frequency noise (LFN), whereas Sco X-1, GX 17+2 and GX 349+2 have small, vertical or absent HBs with a weak or absent HBO and peaked LFN. However, in spite of various details, during *Z*-evolution all these *Z*-sources indicate 6 – 20 Hz N/FBO phenomenon and significant increasing VLFN component at FB. Previous investigations detected no QPO in the FB of GX 340+0 (e.g., Penninx et al. 1991). This fact, along with the absence of type-I X-ray bursts, make one to think about GX 340+0 as a mysterious object. However in this Paper we reveal a typical evolution of GX 340 with FBO events based on our investigation of *RXTE* data set, which helps one to understand the nature of FB in GX 340+0.

In this study we analyze the *BeppoSAX* observations and *RXTE* observations during 1997 – 2009 years for GX 340+0. In §2 we show the list of observations for which we apply our data analysis and the details of X-ray spectral analysis are elaborated in §3. We study how X-ray spectral and timing properties evolve during the different spectral states in §4. We discuss our results and make their comparisons with those known in the literature in §5. We make our final conclusions in §6.

2. Data Selection

We obtain broad band energy spectra of the source using data of three *BeppoSAX* Narrow Field Instruments (NFIs), namely the Low Energy Concentrator Spectrometer (LECS) for 0.3 – 4 keV, the Medium Energy Concentrator Spectrometer (MECS) for 1.8 – 10 keV and the Phoswich Detection System (PDS) for 15 – 200 keV [see Parmar et al. (1997); Boella et al. (1997); Frontera et al. (1997) respectively].

We use the SAXDAS data analysis package for data processing. We make the data spectral analysis in the energy range for which a response matrix of each of the instruments is well determined. We renormalized the LECS data based on the MECS data. We treat relative normalizations of the NFIs as free parameters when proceed with model fitting, but we fix the MECS normalization at a value of 1. We control if these normalizations are in a standard range for each instrument¹. Furthermore, we rebinned the spectra in order to have significant data points. The LECS spectra are rebinned with a binning factor which is not constant over energy (Sect.3.1.6 of Cookbook for the BeppoSAX NFI spectral analysis) implementing rebinning template files in GRPPHA of XSPEC². The PDS spectra are rebinned with a linear binning factor 2, namely we group two bins together leading to the resulting bin width of 1 keV. We apply a systematic error of 1% to all of these spectra. We present a list of the *BeppoSAX* observations implemented in our analysis in Table 1.

We also use publicly available the *RXTE* data sets (Bradt et al. 1993) which were obtained from April 1997 to March 2009. In total, they include 92 observations taken at different states of the source. We apply standard tasks of the LHEASOFT/FTOOLS 5.3 software package for data processing. We use PCA *Standard 2* mode data, collected in the 3 – 20 keV energy range and the most recent release of PCA response calibration (ftool pcarmf v11.1) for spectral analysis. We also apply standard dead time correction to the data. In order to construct broad-band spectra we include the data from HEXTE detectors.

A background corrected in off-source observations is subtracted from the data. We use only data in the 20 – 150 keV energy range in order to avoid the problems related to the HEXTE response and background determination. We use the data which are available through the GSFC public archive (<http://heasarc.gsfc.nasa.gov>). We present a full list of observations covering the source evolution during different spectral state events in Table 2.

We accomplish an analysis of *RXTE* observations spanning twelve years for which 11

¹<http://heasarc.nasa.gov/docs/sax/abc/saxabc/saxabc.html>

²<http://heasarc.gsfc.nasa.gov/FTP/sax/cal/responses/grouping>

intervals indicated by blue rectangles in Figure 1 in Seifina, Titarchuk & Frontera (2013) hereafter STF13. We model the energy spectra using XSPEC astrophysical fitting software and we apply systematic error of 0.5% to all of the analyzed spectra. It is also worth noting that we use public GX 340+0 data from the All-Sky Monitor (ASM/*RXTE*) which show long-term quasi-constancy behavior of mean soft flux during all observation scans.

3. Spectral Analysis

3.1. Choice of the Spectral Model

As a first step we proceed with a model which is a sum of an absorbed thermal component (*Bbody*), a thermal Comptonization component (*Comptb*³) and a *Gaussian* line component. But this model [*single COMPTB* model, $wabs * (Bbody + Comptb + Gauss)$] gave a poor description of about 40% of the data. Significant negative residuals at low energies and greater than 30 keV suggest the presence of additional emission components. For this reason we also attempt the so called *double Bbody* model [$wabs * (Bbody + Bbody + Comptb + Gauss)$], however the best fit of both *RXTE* and *BeppoSAX* spectra has been obtained by implementing the *double COMPTB* model [$wabs * (Bbody + Comptb1 + Comptb2 + Gauss)$] for all available set of the data (see Table 3 for *BeppoSAX* data).

This model describes a scenario in which a Keplerian disk is connected to the *Neutron star* through the transition layer (TL) [see Titarchuk et al. (1998)]. We display our spectral model in Figure 1. We suggest that accretion into a neutron star occurs through the two main regions: an accretion disk [the standard Shakura-Sunyaev disk, see Shakura & Sunyaev (1973)] and TL. Then NS and disk soft photons are upscattered off hot electrons of the transition layer (TL). Thus in our picture, the emergent Comptonization spectrum is formed in TL, where thermal disk and NS seed photons of temperatures kT_{s1} and kT_{s2} upscatter off the TL hot electrons, giving rise to two Comptonized components *Comptb1* and *Comptb2*, respectively. The Earth observer can directly see some fraction of the seed soft photons that justifies an addition of a soft blackbody component of temperature T_{BB} . In Figure 1 we show red and blue photon trajectories which are related to seed and upscattered photons, respectively.

Thus the emergent X-ray spectrum can be presented as a convolution of a soft black body spectrum of normalization N_{com} and color temperature kT_s with a Comptonization

³*Comptb* is an XSPEC contributed model,
see <http://heasarc.gsfc.nasa.gov/docs/software/lheasoft/xanadu/xspec/models/comptb.html>

Green function. Normalization N_{com} is a ratio of the source luminosity to square of the distance D (similar to the ordinary *Bbody* XSPEC model)

$$N_{com} = \left(\frac{L}{10^{39} \text{erg/s}} \right) \left(\frac{10 \text{ kpc}}{D} \right)^2. \quad (6)$$

The applied model *wabs*(Bbody+Comptb1+Comptb2+Gauss)* has the free parameters: the equivalent hydrogen absorption column density N_H ; the energy spectral indices α_1, α_2 ; *seed* color temperatures T_{s1}, T_{s2} ; parameters $\log(A_1)$ and $\log(A_2)$ related to the illumination fractions of the transition layer by disk and NS soft photons respectively, f_1, f_2 , in other words $f = A/(1 + A)$ is the relative weight of the Comptonization component; electron temperatures $T_e^{(1)}$ and $T_e^{(2)}$; normalizations of the BB-like components N_{Com1} and N_{Com2} of the *Comptb1* and *Comptb2* components respectively. To fit the data in the 6 – 8 keV energy range we add to the above model a Gaussian component, which has as parameters: a centroid line energy E_{line} , the line width σ_{line} and its normalization N_{line} .

Note that some of the parameters of the *Comptb* component: $\gamma = 3$ is related to the index of the seed photon blackbody spectrum, $\alpha = \gamma - 1 = 2$, and δ characterizes an efficiency of the bulk inflow effect vs the thermal Comptonization. We neglect this bulk motion effect for NS and we put $\delta = 0$. The bulk inflow Comptonization should take place very close to NS surface but if the radiation pressure from NS surface is high then the bulk motion is suppressed. On the other hand when mass accretion is low then the bulk effect motion effect can be ignored too. In general, for NSs the bulk effect is negligible (compare with a BH case, see ST09).

A value of hydrogen column $N_H = 6.2 \times 10^{22} \text{ cm}^{-2}$ found by Iaria et al. (2006) is used in our model. We also fix the parameter $\log(A_2)$ at a value of 2, on the assumption that NS surface is completely covered by TL (see Fig. 1 for our model geometry).

3.2. *BeppoSAX* data analysis

In Figure ?? we present two representative EF_E spectral diagrams for HB and FB events along Z track of GX 340+0 using *BeppoSAX* data. We show the best-fit spectral diagrams of GX 340+0 using *BeppoSAX* observation 21375002 (*left*) and 20261006 (*right*) on the *top* of this Figure. We present the data by crosses and the best-fit spectral model *wabs*(Blackbody+Comptb1+Comptb2+Gaussian)* by light-blue line. The model components are shown by dark-blue, red, green and crimson lines for *Blackbody*, *Comptb1*, *Comptb2* and *Gaussian*, respectively. *Bottom panels*: $\Delta\chi$ vs photon energy in keV. The best-fit model parameters for HB state (*left* panel) are $\Gamma_1 = 2.04 \pm 0.06$, $kT_e^{(1)} = 19.50 \pm 0.07 \text{ keV}$, $\Gamma_2 = 1.99 \pm 0.02$,

$kT_e^{(2)}=2.76\pm0.09$ keV and $E_{line}=6.80\pm0.06$ keV (reduced $\chi^2=1.28$ for 329 d.o.f), while the best-fit model parameters for FB state (*right* panel) are $\Gamma_1=2.00\pm0.08$, $kT_e^{(1)}=5.42\pm0.03$ keV, $\Gamma_2=2.04\pm0.05$, $kT_e^{(2)}=1.90\pm0.02$ keV and $E_{line}=6.68\pm0.08$ keV (reduced $\chi^2=0.98$ for 319 d.o.f) (see more details in Table 3).

The *Blackbody* temperature kT_{BB} , independently of the source state, is consistent with 0.6 keV (2σ upper limit). The addition of this low temperature blackbody component makes the fit quality noticeably higher. Specifically, the best-fit for HB events (id=21375002) is characterized by reduced χ^2 of 2.1 (331 d.o.f.) for the model *without the low temperature blackbody component*, while χ^2 is 0.99 (329 d.o.f.) for the model *with the low temperature blackbody component*. Moreover the best-fit spectral model for FB events (id=20261006) has χ^2 of 1.7 (321 d.o.f.) for the model without the blackbody component, while χ^2 is 0.98 (319 d.o.f.) for the model with the blackbody component.

For the *BeppoSAX* data (see Tables 1, 3) we obtain that the best fit spectral indices α_1 and α_2 are given by 1.01 ± 0.04 and 0.99 ± 0.05 , respectively, (or $\Gamma_1 = \alpha_1 + 1$ and $\Gamma_2 = \alpha_2 + 1$ are given by 2.01 ± 0.04 and 1.99 ± 0.05 , respectively, for a *double COMPTB* model). We also find that, for the available *BeppoSAX* data, from HB to FB, the seed temperatures kT_{s1} and kT_{s2} change in the ranges 0.7–1 keV and 1–1.7 keV, respectively. We apply a systematic error of 1% to all of the analyzed spectra.

3.3. *RXTE* data analysis

From our analysis of the *BeppoSAX* data we obtain the temperature of the soft *blackbody* component at a value of $kT_{BB} = 0.6$ keV. However *RXTE* detectors cannot give us well calibrated spectra below 3 keV thus in order to fit the *RXTE* data we should fix this *blackbody* temperature at 0.6 keV. We also fix values of kT_{s1} and kT_{s2} at 1.1 keV and 1.5 keV, respectively (see §3.2). in Table 4 we show the best-fit spectral parameters using *RXTE* data. Also we find that the normalization of the *Comptb1* component changes from 2 to 6 in units of L_{37}/D_{10}^2 (where L_{37} is the luminosity of the seed blackbody component in units 10^{37} erg sec $^{-1}$ and D_{10} is distance in units 10 kpc), whereas Γ_1 is around 2 for all observations (see Figs. 2-3). Note that the line width σ_{line} of *Gaussian* does not change much and numerous tests show that $\sigma_{line} = 0.5 - 0.8$ keV.

Furthermore, a detailed analysis XMM-*Newton* spectra of GX 340+0 recently carried out by D’Ai et al. (2009) has revealed that the line profile does not present any strong change for HB. Therefore we fix σ_{line} of Gaussian component at a value 0.6 keV during our fitting procedure while the electron temperature $kT_e^{(1)}$ of *Comptb1* varies from 3 keV to 21

keV (Fig. 4), which is similar to that obtained using the *BeppoSAX* data (see Table 3).

In Figure ?? we show three representative EF_E spectral diagrams for different states along the Z -track. Data are taken from *RXTE* observations 50016-01-01-11 ($Z1$ panel, *Horizontal branch*), 91125-03-01-000 ($Z2$ panel, *Normal branch*) and 50016-01-02-18 ($Z3$ panel, *Flaring branch*). We show the data using black points while the spectral model components are presented by red, green, blue and purple lines for *Comptb1*, *Comptb2*, *Blackbody* and *Gaussian* respectively. One can notice changes of spectral shape for different spectral branches in this Figure. To highlight the effect of variability of different components during evolution between the HB ($Z1$) and FB ($Z3$) states we mark the *Comptb1* component by yellow shaded areas and trace its evolution when $kT_e^{(1)}$ monotonically drops from 21 keV ($Z1$ panel) to 3 keV ($Z3$ panel) (see also Fig. 3 for the electron temperature range).

In Figure 5 three representative EF_E spectral diagrams for different states along Z track (*right panel*) are presented in combination with *color-color* diagram (*left panel*). Data are taken from *RXTE* observations 20053-05-01-01 (*green*, NB) 20053-05-01-02 (*violet*, FB) and 20053-05-01-00 (*red*, HB). On the *right panel* of this Figure one can see changes of spectral shape for different spectral branches which are presented on the *left panel*. Furthermore, the hard emission at 50 – 150 keV (*hard tail*), even if not strongly constrained, appears relatively low along FB, while during HB and NB the *hard tail* becomes stronger.

Our spectral model demonstrates a very good quality for all data sets analyzed in our study. In fact, $\chi_{red}^2 = \chi^2/N_{dof}$, where N_{dof} is a number of degree of freedom, is less than or about 1 for most observations. With high counting statistics for a small fraction of the spectra (less than 2%) χ_{red}^2 can reaches 1.4. On the other hand, χ_{red}^2 is never above a rejection threshold of 1.5 (for 90% confidence level). We should note the energy range, in which we obtain the poor fits (one among 92 spectra with $\chi^2=1.46$ for 67 d.o.f), is associated with the iron line region. Possibly that the iron line feature in the spectrum is more complicated than just one Gaussian line. In fact, for XMM-*Newton* observation D’Ai et al. (2009) detected an asymmetric profile of Fe K_α emission line with an extensive red wing. The authors interpret that this line asymmetry is caused by relativistic smearing in an accretion disk extending close to the NS. However Laurent & Titarchuk (2007) showed that the line asymmetry can be also formed in an optically thick medium expanding with subrelativistic velocities. We must admit that this complex line structure cannot be resolved by the *RXTE* data.

We should also point out some differences between our best-fit values of the photon index Γ and those found by us in the literature. In particular, the photon index Γ , estimated by Iaria et al. (2006) for *BeppoSAX* observations ($S3 - S6$) gives the values of Γ is about 2.5 whereas Γ is about 1.7 for *RXTE* observations ($S5$) [see for details Church et al. (2006)]. However, this difference of Γ values is a result of using different spectral models. For example,

Church et al. used an additive model of a blackbody, a power law components and a single Comptonized component (*Comptt*).

Thus using broad band *BeppoSAX* observations we obtain our spectral model parameters and we can evaluate the total spectral evolution of GX 340+0 from 3 to 150 keV due to the elongated observations by the *RXTE*.

4. Overall pattern of X-ray properties

4.1. Color-color diagrams

To study the properties of GX 340+0 during different spectral states we make use of hard colors (HCs) and soft colors (SCs) to demonstrate different configurations in HC versus SC. Namely, in Figure 9 in STF13 we collected color-color diagrams (CCDs) (*left column*) and hardness-intensity diagrams (HID) (*right column*) of GX 340+0. The ordinate and abscissa of color-color diagrams show the flux ratios: (a) [16-50 keV/3-16 keV] and [3-5 keV/7-10 keV]; (c) [20-50 keV/3-10 keV] and [10-20 keV/3-10 keV]; (e) [10-20 keV/3-10 keV] and [9-16 keV/3-5 keV] while hardness-intensity diagrams demonstrate flux ratios: (b) [16-50 keV/3-16 keV]; (d) [20-50 keV/3-10 keV]; (f) [7-10 keV/3-5 keV] versus flux (3-50 keV) measured in units of 10^{-9} erg/s/cm². In this way, one can observe different shapes of the *Z* pattern of GX 340+0. For the presented observations (see Table 1 and 2) we found the source in three spectral states: the flaring branch (FB), the normal branch (NB) and the horizontal branch (HB).

The identification of CCD states was made using simultaneous timing and spectral analysis, see §4.3 and we have revised the previous similar investigation obtained using EXOSAT and *RXTE* data, see Penninx et al., 1991; Jonker et al., 1998, 2000, respectively (see §3). Both the color–color and color–flux behaviors depend on the photon energy bands in which either color or flux are evaluated. In particular plots, see panels (a), (b) and (d), there is evidence of HB, while in other ones this branch is not apparent. The evidence of FB appears only from panels (a) and (b). Instead, independently of the energy band, NB is always apparent. A comparison of the color–flux behavior of different compact X–ray binaries that host NS or a BH is shown in Fig. 6.

It should be noted that FB flux behavior of GX 340+0 corresponds to dips in the light curve (see Figs. 11-12 in STF13). Indeed, when the source intensity dips, both the soft and hard colors increase (see panel (a) in Fig. 9 in STF13). This behavior is completely different from FB one observed in Sco X-1, GX 17+2 and GX 349+2, while it is similar to the FB behavior seen in Cyg X-2. A discussion of this phenomenon is given by Hasinger et al. (1989)

and Kuulkers & van der Klis (1996).

4.2. Evolution of X-ray spectral properties in different spectral states

We find properties of the HB-NB-FB state of GX 340+0 based on their spectral evolution in the 3–150 keV energy range using *RXTE* data.

The general picture of HB-NB-FB states is shown in Figure 5 where we put together the HB, NB and FB spectra to display the GX 340+0 spectral evolution (see also panels for HB and NB spectra in Fig. ??). One should point out different spectral shapes related to the different spectral states. Figures 3 and 7 in STF13. illustrate the model components while Figs 11–12 in STF13 demonstrate the evolution of these components. As we indicate below (see §4.2.1), using of the best-fit model, in all spectra independently of the state a strong Comptonized component (*Comptb2*) related to the upscattering of the NS seed photons off the TL innermost part is dominant (see illustration of this effect in Fig. 1).

In these two states, the Comptonization component, *Comptb1*, related to upscattering of the disk seed photons by TL electrons of the temperature in the range of 8–21 keV, extends up to 150 keV. Instead, in FB spectra (Fig. 7 in STF13 panel for the FB spectrum) the Comptonized component *Comptb1* is characterized by a decrease of the electron temperature to 3 – 6 keV and thus the spectrum only extends to 50 keV. Generally, the *Blackbody* and *Comptb1* components are relatively weak with respect to *Comptb2*. We find that values for the electron temperature $kT_e^{(2)}$ is close to that of $kT_e^{(1)}$, only during *flaring branch*, when $kT_e^{(1)}$ monotonically decreases to 3 keV, while $kT_e^{(2)}$ shows low variability about 3 keV during all spectral states. HC ratio [10-50 keV]/[3-10 keV] drops from 0.3 to 0.1 during HB-FB spectral evolution (see Fig. 4).

Thus using Figure 4, one can *find* one-to-one correspondence between spectral states of GX 340+0 and value of $kT_e^{(1)}$ within the limits, marked by horizontal blue dashed lines. The electron temperature $kT_e^{(1)}$ varies by a factor of 10 during these states while Γ_1 and Γ_2 , for each of these spectra, only vary from 1.9 to 2.1 (see Fig. 3).

Using χ^2 -statistic criterion, we test the hypothesis that the approximate value Γ_{appr} of both indices Γ_1 and Γ_2 is $\Gamma_{appr} \approx 2$. We calculate the distribution of $\chi_{red}^2(\Gamma_{appr}) = \frac{1}{N} \sum_{i=1}^N \left(\frac{\Gamma_i - \Gamma_{appr}}{\Delta\Gamma_i} \right)^2$ versus of Γ_{appr} . We find a sharp minimum of the function $\chi_{red}^2(\Gamma_{appr})$ of about 1 for two Comptonized components (*Comptb1* and *Comptb2*). Namely $(\Gamma_1/\Gamma_2)_{appr} = 1.99 \pm 0.01/2.00 \pm 0.01$ with a confidence level of 67% and $(\Gamma_1/\Gamma_2)_{appr} = 1.99 \pm 0.02/2.00 \pm 0.02$ with a confidence level 90% for 92 d.o.f. (see also Figure of $\chi_{red}^2(\Gamma_{appr})$ for 4U 1728-34 in ST11). We should point out that the indices Γ_1 , Γ_2 are also independent of the

normalizations of *Comptb1*, *Comptb2*; L_{39}/d_{10}^2 and $kT_e^{(1)}$, $kT_e^{(2)}$ (see Figs. 2–3). Relying on *BeppoSAX* observations and their analysis FT11 propose that Γ varies about 2 for many NSs which are seen from the hard to soft states. FT11 relate the spectral state to a value of electron temperature T_e and they demonstrate that the photon index $\Gamma = 2 \pm 0.2$ (or the spectral index $\alpha = 1 \pm 0.2$) while kT_e varies in the range from 2 to 25 keV.

4.2.1. Light curves and related spectral characteristics

We find a number of X-ray flaring events of GX 340+0 using *RXTE* data obtained during 2000–2001 (*R5*, *R6* sets) with a full rise-decay evolution. We examine generic spectral and timing properties which can be found during these spectral state events. In Figures 11 in STF13 we show the details of our spectral analysis of the *RXTE* observations applying our model $wabs * (bbody + Comptb1 + Comptb2 + Gaussian)$.

In Figure 11 in STF13 we show the changes of count rate (2-9 keV) in counts s^{-1} with 16 s time resolution, the *hardness ratio* coefficient HC [10-50 keV]/[3-10 keV], the electron temperatures $kT_e^{(1)}$ (*red*)/ $kT_e^{(2)}$ (*blue*) of *Comptb1*/*Comptb2* components, the normalizations of *Comptb1* and *Comptb2* components (*red* and *blue* respectively) and the index $\alpha = \Gamma - 1$ for *Comptb1* and *Comptb2* components (*red* and *blue* points respectively) during the 2000 events (*R5* set) of time period from MJD 51722 to MJD 51777. The dipping phases of the light curve, which are related to the *flaring branch* (based on timing analysis, see also Fig. 9 and § 4.3) are marked with blue vertical strips. Note that during these dipping phases of the light curve the *hardness ratio* coefficient (HC) is usually less than 0.15.

Additional flaring events are presented in Fig. 12 (STF13) where we display details similar to those shown in Fig. 11 in STF13 but for 2001 events (*R6* set) during the MJD 51920–MJD 51925 period. Note the temperature $kT_e^{(1)}$ also correlates with the *hardness ratio* coefficient HC [10-50 keV]/[3-10 keV] and it monotonically diminishes from 20 keV to 3 keV during the HB-FB transition.

It is clear to see, using Figs. 11- in STF13 that the electron temperature $kT_e^{(1)}$ (*red* points) steadily decreases from 20 keV during the transition from HB to NB, reaching its minimum at 3 keV in flaring FB [MJD 51774 – 51774.3, 51776.5 – 51776.7 (Fig. 11 in STF13); MJD 51921.4 – 51921.7, 51922.4 – 51922.7 (Fig. 12 in STF13)] while the electron temperature $kT_e^{(2)}$ (*blue* points) is almost constant at the mean value of about 3 keV. On the other hand, the *Comptb* normalizations N_{Com1} and N_{Com2} are weakly correlated with the variations of *hardness ratio* coefficient HC and count rate in 2-9 keV energy band, with the X-ray emission related to *neutron star* that dominates during all spectral states (see Fig. 7).

The normalization of $Comptb2$, N_{Com2} is low variable around its mean level of $0.15 \times L_{39}/D_{10}^2$. The X-ray contribution from the Comptonization of the disk seed photons is weaker by factor 2 than that related to the NS seed photons. during all spectral states (Fig. 7). We should point out that the indices α , shown in the bottom panels of Figures 11-12 in STF13 only slightly vary with time around 1 (or the corresponding Γ varies around 2).

4.2.2. Comptonized emission

Note that the Comptonization fraction f_1 changes from 0 to 0.5 as shown on Fig. 8 (*red triangles*). Thus, in most cases of HB and NB states, the soft radiation of GX 340+0 dominates and is only slightly modified by reprocessing in the TL ($f_1 = 0.1 - 0.2$). Then, during the transition to FB, f_1 monotonically increases from 0.2 to 0.5 when $kT_e^{(1)}$ decreases from 6 keV to 3 keV. Consequently, the Earth observer directly sees a smaller fraction of disk emission ($1 - f_1$) in FB. We remind a reader that the Comptonization fraction f_2 , related to the NS surface seed photons upscattered off the TL electrons, is about 1. We freeze $f_2 = 1$ on the assumption that NS is completely surrounded by its TL (see Fig. 1 for the model geometry).

4.3. Timing properties during HB-NB-FB evolution

We analyze the *RXTE* light curves using the *powspec* task from FTOOLS 5.1. We perform the timing analysis *RXTE*/PCA data using the *event* mode in the 13 – 30 keV range. The event mode time resolution is 1.2×10^{-4} s. We obtained PDSs in the 0.1 – 500 Hz frequency range with 0.001-second resolution. The contribution due to Poissonian statistics and Very Large Event Window for all PDSs were subtracted.

We demonstrate an evolution of the 3 – 150 keV flux during the 2000 (*R6*) events (see *top panel* of Figure 9). Here *red/blue* points A, B, and C mark moments at MJD = 51773.2/51774.7, 51773.6/51774.1 and 51772.2/51774.6 related to different phases of *Z*-state evolution. In addition, *green* points D_1/D_2 indicate moments at MJD=51774.27/51776.68 of FB where we find ~ 6 Hz QPO at *flaring* branch of GX 340+0 (see also Fig. 10 and details below).

We present PDSs for 15-30 keV range (*left column*) plotted along with spectral diagrams $E * F(E)$ (*right column*) for A (*red*, top), B (*blue*, middle) and C (*blue*, bottom) points of the light curve in the *bottom panel*. A typical exposure time for each PDS is 3-4 ksec. In particular, the accumulation time for PDS during 50016-01-02-16 observation (see Fig. 10)

is 3876 sec.

An evolution of timing properties over the Z track is clearly seen in this segment of the light curve (see *top panel* of Fig. 9). Strong oscillations with centroid frequency at 50 Hz are observed at HB (see Fig. 9, left panel, C *red* PDS related to MJD=51772.2). While broad oscillations ranging in 2 – 10 Hz (B *red*, 50016-01-01-06, MJD=51773.6; C *blue*, 50016-01-02-04, MJD=51774.6) and ~ 50 Hz (A *blue*, 50016-01-02-12, MJD=51774.7) are detected during middle and hard apex of *normal* branch (NB), respectively. Relatively strong noise component (LFN) is present (up to 10% rms) and NB (up to 2% rms) but none of these features is seen during FB when a very low frequency noise (VLFN) component arises (B *blue*, 50016-01-02-18, MJD=51774.1). We show the $E * F(E)$ diagrams (A2, B2, C2 panels) corresponding to the related PDSs (A1, B1, C1 panels) on the right. We present the data by black points whereas we demonstrate the spectral components using dashed red, green, blue and purple lines for *Comptb1*, *Comptb2*, *Blackbody* and *Gaussian* respectively.

We also reveal ~ 6 Hz QPOs in FB during two “dipping/flaring” events ($R5$ data set), which are not detected previously in GX 340+0, but present in power spectra of all other Z sources during this state. More specifically, during first dipping phase of light curve [for example, see point B *blue* (MJD 51774.08) in Fig. 9] we find only strong VLFN component (see *blue* histogram, in the middle panel of left column) without QPO feature. While during the last part (*exit phase*) of the intensity dip we find a broad QPO Lorentzian centered at 6 Hz (FWHM= 11.7 ± 4.5 Hz, rms= $6.3 \pm 1.0\%$) in addition to VLFN (rms= $4.1\% \pm 0.4\%$, $\alpha_{LF}=1.4 \pm 0.3$, $\chi^2 = 139$ for 102 d.o.f; we present all parameter errors corresponding to 1σ confidence level). The related point of the light curve is indicated by *green* point D_1 observed on August 18, 2000 (50016-01-02-16, MJD=51774.27) seen in Fig. 9. We make the corresponding $\nu \times power$ diagram for this event in 0.01 – 150 Hz range (see *left* panel of Fig. 10). We should point out that the energy spectrum at point D_1 (*right* panel of Fig. 10) is a typical FB spectrum [e.g., see also Z3 panel of Fig. 7 in STF13 with a relatively low contribution of the hard emission at 70– 150 keV (hard tail), while at HB and NB the hard tail becomes stronger (*red/blue* points A and C in Fig. 9)].

During another FB event (see *green* point D_2) we detected the same behavior of PDS, characterized by the absence of QPO at the beginning and middle of FB but the presence of broad centered QPO at frequency about ~ 6 Hz in the *bottom phase* of FB (50016-01-02-01, MJD=51776.68) in addition to strong VLFN component. Note that the energy spectra for both points (D_1 , D_2) are very similar (their best-fit parameters are mostly identical).

Note that the energy spectra of $D_{1/2}$ points are similar to that indicated by B (blue) point (and generally typical to FB spectra, see Z3 diagram of Fig. 7 in STF13 but the related power spectra are completely different. In fact, we reveal a broad QPO, centered at

$\nu_{FBO} \sim 6$ Hz, for the first time in the PDS of GX340+0 (see Fig.10).

Thus, we find that the evolution of spectral and timing characteristics is consistent with previous analysis of GX 340+0 using EXOSAT, *Ginga*, *RXTE* and XMM-*Newton* data [see e.g. Penninx et al. (1991), Jonker et al. (2000)]. In addition to known timing features of the *Z* pattern of GX 340+0 we reveal ~ 6 Hz FBO, which was not found previously, while it is usually observed in other *Z* sources such as Sco X-1 and GX 17+2. This kind of QPOs are seen in the lower FB, and also seen in NB of those *Z*-sources. The frequency starts to increase from $\nu_{FB} \sim 6$ Hz to $\nu_{NB} \sim 20$ Hz when these particular sources progress from FB to NB. In the lower FB, this QPO frequency ~ 6 Hz becomes very broad and then vanishes in the noise.

Follow to Kuulkers & van der Klis (1996) and Titarchuk et al. (1998), we can interpret the appearance of this QPO phenomenon during FB in terms of the *Transition Layer* (TL) model involving a radial flow and a thick, torus-like CC (see illustration of this interpretation in Fig. 11). To conclude the above analysis, we should point out once again that the *identification* of CCD states of GX 340+0 along *Z*-track is made using timing (presented here) along with spectral analysis (see §3).

4.4. Comparison of spectral and timing characteristics of *Z* source GX 340+0 and *atoll* sources GX 3+1 and 4U 1728-34

In this Paper, we study the correlations between spectral, timing characteristics, and $\dot{M}$ seen in X-ray spectral range for GX 340+0 during its evolution across HB-NB-FB track in order to investigate a common behavior of *Z* and *atoll* sources. In this way we proceed with a comparative analysis for three sources: *Z*-source GX 340+0 and *atolls*, GX 3+1 and 4U 1728-34 using the same spectral model which consists of low temperature *Blackbody*, *Comptonized* continuum and *Gaussian* line components.

4.4.1. Quasi-constancy of the photon index

Z-source GX 340+0 and *atolls*, GX 3+1, 4U 1728-34 show an identical behavior of Γ vs $\dot{M}$ (or COMPTB normalization), namely the index Γ is strongly concentrated around 2 (see Fig. 2 for that in GX 340+0). According to FT11, ST11 and ST12, this result can point out that the cooling flux of the soft disk photons is much less than the gravitational energy release in the *transition layer* for all of these three sources.

4.4.2. The difference and similarity of the electron temperature kT_e ranges in GX 340+0, GX 3+1 and 4U 1728-34

One can see, using Figure 8, the ranges of the TL electron temperature kT_e for state evolutions of these three sources. As clearly seen in this Figure (see also Fig. 12) kT_e of *Z*-source GX 340+0 ranges from 3 to 21 keV, while that of *atoll* 4U 1728-34 is slightly more narrow, from 3 to 15 keV and kT_e of *atoll* GX 3+1 just varies around 3 keV. Note that all objects have a common temperature interval, 3 – 4 keV.

As it is shown in Table 5 variations of of *blackbody*, and seed (NS and disk) photon temperatures are comparable for all of these three sources: $kT_{BB} \simeq 0.6$ keV and $kT_s = 1.1 - 1.7$ keV, respectively. But variations of kT_e are different. This difference of kT_e – ranges for all of these three sources is quite evident. Whereas GX 340+0 and 4U 1728-34 show a complete cycle of state evolution, namely, HB-NB-FB track for GX 340+0 and 4U 1728-34 evolves from the extreme island state (EIS) to the upper banana (UB) state (see Di Salvo et al. 2001; ST11), GX 3+1 demonstrates only a short LB (lower banana)-UB track on the CCD. It is clear from Figure 8 which shows the track of GX 3+1 just for a part of the full track [see definition of a state sequence in Hasinger & van der Klis (1989)].

But GX 340+0 shows almost the same kT_e range as that in *atoll* 4U 1728-34, while it demonstrates a different timing evolution. How different *Z* and *atoll* sources are one can clearly see from Fig. 12, which shows the luminosity in terms $N_{Com} \times D^2_{10}$ versus T_e . *Z*-source GX 340+0 (*pink* points) is much brighter than *atolls*, 4U 1728-34 (*blue* points) and GX 3+1 (*red* points).

4.4.3. Comparison of spectral evolution as a function of the luminosity for *Z* source GX 340+0 and *atoll* sources GX 3+1 and 4U 1728-34

Now we can make a comparative analysis of spectral parameter evolution for GX 340+0 and GX 3+1, 4U 1728-34 based on *luminosity* of soft disk photons, which is proportional to disk mass accretion rate and inversely proportional to square of distances to these sources (see Eq. 6 and Table 5). We should remind a reader that *atoll* sources 4U 1728-34 and GX 3+1 demonstrate only one Comptonized (*Comptb*) component in their X-ray spectra [Seifina & Titarchuk (2011), Seifina & Titarchuk (2012)], while *Z*-source GX 340+0 reveals two Comptonized components (*Comptb1* and *Comptb2*). The first *Comptb1* component is formed by up-scattering processes in TL for seed photons of temperature $T_{s1} \leq 1$ keV coming from the disk, while the second *Comptb2* component is formed as a result of up-scattering processes in TL for seed photons of temperature $T_{s2} \sim 1.5$ keV coming from the neutron

star. In contrast with GX 340+0, the spectra of *atolls*, 4U 1728-34 and GX 3+1 contain only one Comptonized (*Comptb*) component, which is related to up-scattering processes in TL for seed photons of temperature $T_s \lesssim 1$ keV, presumably coming from the disk. Therefore we should compare spectral components of the same origin for considered *Z* and *atoll* sources. Namely, *Comptb1* component (with f_1 illumination parameter) for GX 340+0 and *Comptb* components (with f illumination parameter) for 4U 1728-34 and GX 3+1.

Specifically, for GX 340+0 the distance is about 10 kpc (Fender & Henry 2000), whereas for 4U 1728-34 and GX 3+1 the distances are 4.5 kpc and 4.2–6.4 kpc, respectively [see van Paradijs (1978) and Kuulkers & van der Klis (2000)]

We present the *luminosities* of these sources as a function of kT_e in Fig. 12. For GX 340+0 (*pink*), we show a variable N_{Com1} normalization only. GX 340+0 demonstrates much higher luminosities by factor 5 than that in 4U 1728-34 and GX 3+1 and kT_e in GX 340+0 widely varies, from 3 to 21 keV. According to FT11, ST11 and the presented study, kT_e for *atoll* and *Z*-sources varies from 2.5 to 25 keV. Specifically, a common range of kT_e for GX 340+0, GX 3+1 and 4U 1728-34 is around 3 keV and the low limit of kT_e around 2.5 keV occurs at the peak luminosity for 4U 1728-34 (see ST11) and during a rise of luminosity for GX 3+1 and during FB stage for GX 340+0 (see Figs. 8, 12).

We should note that common states in L vs T_e diagram for all these sources take place during rapid transitions in FB and LB – UB for GX 340+0 and GX 3+1, 4U 1728-34, respectively. These states can be possibly traced within a narrow kT_e interval from 2.5 to 4.5 keV.

4.4.4. The difference and similarity of time scales for state evolution for GX 340+0, GX 3+1 and 4U 1728-34

We point out that all these three objects show a full range evolution track but in different time scales. Specifically, *Z*-source GX 340+0 and *atoll* GX 3+1 demonstrate electron temperature and *luminosity* variations occur on the time scales of hours-days, while *atoll* 4U 1728-34 show a full range of the model parameter variations on significantly larger time scale (days-months). However, the low electron temperature phase of *atoll* 4U 1728-34 occurs at the same short time scale as that for GX 340+0 and GX 3+1.

4.4.5. Correlation of illumination parameter f with electron temperature kT_e and its relation with different states in the color-color diagram

Using Table 5 one can notice that the variation of Comptonization fraction f is larger for GX 3+1 ($0.2 - 0.9$) than that for 4U 1728-34 ($0.5 - 1$) and GX 340+0 ($0.01 - 0.5$). Thus it is obvious that illuminations of the Transition Layer by soft photons are different for these three sources.

In Figure 8 we plot the electron temperature T_e versus Comptonized fraction f for Z-source GX 340+0, and *atolls*, GX 3+1 and 4U 1728-34 during different spectral states. *Red*, *green* and *blue* points correspond to *RXTE* observations of GX 340+0, GX 3+1 and 4U 1728-34 respectively. On the left hand side of the Figure one can see a sequence of CCD states HB, NB, FB (the horizontal, normal and flaring branches respectively) which are presented according to the standard Z-scheme (Hasinger et al. 1989). Based on the obvious correspondence between kT_e and spectral states (Fig. 4) and timing evolution of GX 340+0, kT_e is directly related to the sequence of CCD states in GX 340+0. Note that recently ST12 revealed one-to-one correspondence between spectral states, kT_e and f for *atolls*, GX 3+1 and 4U 1728-34. The direction in which the inferred $\dot{M}$ increases is indicated by arrows.

In Fig. 8 one can see three different tracks on the plot of kT_e versus f for GX 340+0, GX 3+1 and 4U 1728-34 and how these tracks are related to the standard sequence of CCD states. When f rises, kT_e monotonically decreases from 21 keV to ~ 3 keV for GX 340+0, and from 4.5 keV to ~ 2.3 keV for GX 3+1, while 4U 1728-34 shows a more complex example. At high temperature state (EIS), when kT_e decreases, f slightly changes from 0.9 to 1. Further as kT_e drops from 12 keV to 4 keV, f decreases from 0.9 to 0.5. Ultimately, at the low-temperature state (LB-UB) f surges from 0.5 to 1. Thus, one can see that the CCD state evolution can be observable as a relation between kT_e and f too.

4.5. Comparison of spectral hardness diagrams for *atoll*, Z and BHC binaries

To compare evolution properties of GX 340+0 with *atolls* and BHC sources in terms of flux (or luminosity), we plot HC (10-50 keV/3-50 keV) versus the flux in the 3-50 keV range to create our hardness-intensity diagram (HID) for four sources (see Fig. 6): Z-source GX 340+0 (*red*), *atolls*, GX 3+1 (*green*) and 4U 1728-34 (*blue*), and BHC SS 433 (*crimson*). We should remind a reader that Seifina & Titarchuk (2010), hereafter ST10, found a correlation of the index with mass accretion rate $\dot{M}$ and a clear signature of the index saturation when $\dot{M}$ increases. Thus ST10 argue that SS 433 should be a black hole because the index changes and saturates with $\dot{M}$ in BHs only.

As it appears from Figure 6, NS (*atoll* and *Z*) binaries trace a specific elongated or diagonal track in this HID over a larger range of luminosity and (hard) colors than that of BHC SS 433. However, *atoll* sources systematically are fainter than *Z* and BHC sources and show much harder spectra when they are in the low state [see e.g. van der Klis (1994a) for details of comparison of NS and BH sources]. In fact, when *atolls* in the soft state, their spectra are pretty similar to the spectra of *Z*-sources. However *Z*-sources are usually softer than *atolls* but not in the case of *GX 340+0* which is an *intermediate* case between *atolls* and BHs in terms of their luminosities. This fact possibly reflects *intermediate* rate of mass transfer in *GX 340+0* among *atoll* and BH sources [see also a review by van der Klis (1994)].

The resulting luminosity in *GX 340+0* is in the range of $(0.3 - 0.7) L_{\text{Edd}}$. This estimate can be derived using the NS seed photon temperature which varies in the 1-1.7 keV range (see Table 3). Type-I X-ray bursts are not observed in *GX 340+0*. This observational fact can be a consequence of high $\dot{M}$ in this source. Usually type-I X-ray bursts are seen in low-hard states of NS when the resulting radiation pressure between outbursts is quite low. Thus nothing can stop to accumulate plasma in the NS surface until the plasma column reaches its critical weight which leads to the explosion (X-ray burst). Thus a certain level of the surface weight should be attained in order to start an X-ray burst. Narayan & Heyl (2002) demonstrated that X-ray bursts (instability) can occur for a wide range of the temperature of the accreted flow T_{in} (and corresponding emergent luminosity L). Although this instability is less probable for higher T_{in} . Probably this is precisely our case of *GX 340+0* where L is order of Eddington or little bit less ($L \gtrsim 0.5 L_{\text{Edd}}$).

5. Discussion

To further study the similarities between *Z* and *atoll* sources we have investigate how X-ray spectral, timing characteristics and mass accretion rate $\dot{M}$ are related in *Z*-source *GX 340+0* during the evolution across HB-NB-FB track. This investigation helps us to constrain the models related to X-ray spectral formation in compact objects.

5.1. Does mass accretion rate monotonically increase during HB→NB→FB evolution?

We observe a quasi-constancy of N_{Com2} during HB-NB-FB evolution, which can be presented as an argument against the commonly adopted view that $\dot{M}$ monotonically increases along the *Z*-track, since $\dot{M}$ is related to the normalization of the Comptonized component.

In fact, N_{com} (see Eq. 6) is proportional to the soft photon luminosity which is its turn proportional to $\dot{M}$.

We detect spectral and timing evolution of GX 340+0 similar to that of *dipping* Z-source Cyg X-2, where we see that the spectral features of the source are strongly influenced by its high inclination angle (Vrtilek et al., 1999, Titarchuk et al. 2007). Cowley et al. (1979) reported that Cyg X-2 is observed at an inclination between $65^\circ - 75^\circ$. Farinelli et al. (2009) analyzing *BeppoSAX* observations of Cyg X-2 and using a sum of two COMPTB components, successfully described the spectral behavior of the source from its HB to NB. Consequently we can suggest that modulation of the flux from the TL outer part and TL innermost part (related to N_{Com1} and N_{Com2} , respectively) is a result of geometric effects.

In fact, there is no direct information on the binary inclination angle i of GX 340+0, based on radial velocity analysis. The only indication on inclination angle i of 35° for GX 340+0 is deduced using the so called “relativistically smeared out” profile of Fe K_α line. D’Ai et al. (2009) suggest that the iron line profile can be produced as a result of reflection of X-ray hard radiation, forming in Comptonization region, from relatively cold NS and disk surfaces. But the spectral features due to reflection should be seen only if the spectral index of the incident flux onto NS or disk surfaces is essentially less than 1 (Laurent & Titarchuk 2007) not like in our case of GX 340+0 (see Figs. 2-3).

An assumption of high inclination angle of GX 340+0 leads us to suggest that $\dot{M}$ can be greater at FB, when we can possibly see only some relatively small part of X-ray emission from NS surface due to Comptonization region screening effect (see Fig. 11 for explanation of geometrical constraints).

Furthermore, in the case of low inclination angle we would also see strong BB component and monotonic increase of the normalization of Comptonized component along HB – FB track. However, our spectral analysis indicates low BB contribution during all Z-states (see Sect. 4.2.1). This fact can be considered as an additional argument in favor of high inclination angle for binary GX 340+0 (see discussion in Sect. 5.3). In turn, the influence of high inclination on the soft X-ray flux can result in some distortion of real *normalization* value. Thus, the normalization parameter N_{com} is not a reliable indicator for $\dot{M}$ in the GX 340+0 case.

5.2. Detection of QPOs in the Flaring Branch of GX 340+0

In this study we report a discovery of 6 Hz QPO in the FB of GX 340+0. Specifically, when the source enters the upper part of FB, a QPO peak appears with a central frequency

of about 6 Hz. This part of FB corresponds to a dip (more specifically, the final part of the dip) in the light curve. Note that these QPOs exhibit a number of clear differences with respect to FBOs that were observed in Sco X-1 and GX 17+2. In these two sources, there is a continuous evolution between the NB and FB and a value of the QPO frequency steeply rises near the NB-FB state evolution, but never disappears (see Dieters & van der Klis 2000, Penninx et al. 1990). But in our data the 6 Hz QPO is clearly seen in NB (which becomes broad in lower FB), but there is no QPO in the beginning and middle of FB.

The QPO frequency seen in Cyg X-2 is 26 Hz, is the highest ever seen in FB of any Z-source, with a FWHM of only 3 Hz (Wijnands et al 1998). In Sco X-1 and GX 17+2 FBOs can be observed up to 20 Hz as the source moves along FB, but at that point the QPO usually becomes very broad (with the width of ~ 10 Hz, see e.g. Dieters & van der Klis 2000), and further up in the FB, this QPO drops into the background noise components (Middleditch & Friedhorsky 1986). The FBOs, which are seen before, have been related to a hard energy spectrum, whereas the QPOs that we reveal, are associated with a soft spectrum. We should note that Hasinger et al. (1990) reported some excess of power between 5 and 20 Hz in GX 340+0 during a dip in FB, which they interpreted as a possible FB QPO. In view of the differences in the FB properties, we suggest that the 6 Hz QPO which we see in FB, may be caused by a similar mechanism as the usual N/FBO. It is remarkable that the QPO disappears when the source enters into the dip and appears once again when GX 340 leaves the dip. This can suggest that during the dip the emergent emission is suppressed because the line of sight to the source is affected by an absorbing or scattering medium. Fortner et al. (1989) suggest that NBO arises due to a feed-back loop between the radiation force and the accretion rate in an approximately radial inflow fed by matter that was initially in the accretion disk, but has lost its angular momentum due to photon drag. In this model, FBOs are also assumed to originate as a result of oscillations in this radial flow, at super-Eddington accretion rates.

In this way, as it has been figured out by a number of authors (e.g. van der Klis et al. 1987, Alpar et al. 1992 and Titarchuk & Osherovich 2001,), near the Eddington limit the radiation-pressure dominated region of the disk might be puffed up which for certain inclinations could partially obscure the innermost part of the X-ray emitting regions, and their oscillations.

5.3. Possible indication of a high inclination of binary GX 340+0

The fact that the intensity shows a dip, when the QPO appears, can be an indication that the torus enters into the line of sight. Adopting a torus-like shape for the innermost part

of the source when the source becomes near-Eddington luminosity regime we can explain the differences in the shape of FB in HID between different Z -sources and the correlated differences that are observed in N/FBO properties (see Kuulkers et al. 1994). All these differences can be due to differences in binary inclination angles (Fig. 11). We suggest that the sequence of events, that is observed, takes place when the source approaches the Eddington limit. It is possible that a gradual increase in the geometrical thickness of the inner region (Compton cloud) is due to a gradual increase of total $\dot{M}$ (Fortner et al. 1989). What the observer sees strongly depends on the binary inclination angle i . At low i , the torus never comes to the line of sight and thus during FB the X-ray intensity (or flux) always increases, the N/FBO are seen all the way between 6 and 20 Hz (e.g. Sco X-1, GX 17+2). But at higher inclinations i , FB is seen out by the Earth observer when the flux increases and thus the Compton cloud (torus) shielding the NS enters into the line of sight, similar to that in case of Cyg X-2 (see illustration of this event in Fig. 11).

In general, sources observed at angles $> 65^\circ$, the so-called dipping sources, show regular dipping, modulated at the orbital period by the presence of an extended bulge. The observational situation can be different for GX 340+0 where achromatic dipping seems to relate to some turbulent accretion episodes at the inner edge of the disk. For this geometry, one can argue that inclination should not be so important. However if the inclination angles i are quite small one can see the direct BB component of the NS while in the case of GX 340+0 this NS BB component is not observed and it can be an additional argument that the Earth observer sees GX 340+0 at relatively high inclination angles.

In fact, if the transition layer (or Comptonization region) is puffed up then the NS BB direct radiation is not seen and in this case there is no dependence on the inclination angle. But if the luminosity less than Eddington then there is a difference between the low and high inclination cases. At the former case one can see the direct BB component of the NS while in the latter case the BB component is not seen. The NS BB radiation is up-scattered off Compton cloud plasma (the TL innermost part). This can be our case, because independently of the mass accretion rate (or spectral state), the illumination factor f_2 of the NS soft photons is always about 1 (see Sect. 4.2.1). It means that we always see this source at high inclination angle independently of $\dot{M}$.

6. Conclusions

In this Paper, we demonstrate the results of our study of spectral-timing correlations and their dependences on mass accretion rate observed in GX 340+0 with the *RXTE* and *BeppoSAX*. We study the similarities between Z and *atoll* sources to make constraints and

figure out models which can explain X-ray spectral formation in these compact objects. In our study we use the broad energy band and adequate spectral resolution of the *BeppoSAX* detectors in the 0.1–200 keV range combined with the *extensive RXTE* observations in the 3–150 keV range.

We find that the X-ray broad-band energy spectra of GX 340+0 can be properly reproduced by an additive model, composed of low-temperature *Blackbody* component, two Comptonized components with different *seed* photon temperature ($T_{s1/s2}=1/1.5$ keV). We should also add the iron-line (*Gaussian*) component to fit the *BeppoSAX* and *RXTE* data of GX 340+0. Spectral analysis using our model demonstrates that the photon indices Γ_1 and Γ_2 of these Comptonized components only slightly vary around 2, namely $\Gamma_1=1.99\pm0.02$ and $\Gamma_2=2.00\pm0.02$. While the electron temperature $kT_e^{(1)}$ of the outer part of the transition layer (TL) changes from 21 to 3 keV and the electron temperature of the TL innermost part $kT_e^{(2)}$ varies around 3 keV along *Z*-track from HB to FB.

Based on *RXTE* data we detected ~ 6 Hz quasi-periodic oscillations at the *flaring branch* (FB), which were not detected previously in GX 340+0, but this type of QPO frequency have been revealed in PDSs of all other *Z*-sources at FB.

We explain this *stability* of the indices Γ of Comptonized components around a value of 2. The index stability now found for Comptonized spectral components of the *Z* source GX 340+0 during the state evolution of the source is similar to that previously established in *atolls*, 4U 1728-34, GX 3+1 and shown for quite a few of other LMXBs (see Farinelli & Titarchuk, 2011).

We also find in the *BeppoSAX* spectra that there are two blackbody components, one, with relatively low color temperatures 0.6-1 keV, is probably related to the disk emission and another one of color temperatures about 1.5 keV probably corresponds to the NS surface emission.

We establish that kT_e monotonically arises in the 3–21 keV range when GX 340+0 evolves from FB to HB (see Fig. 8).

In ST11 we show that if the energy release Q_{cor} in the Transition Layer is much higher than that in the disk Q_{disk} i.e. $Q_{cor}/Q_{disk} \gg 1$, then the resulting photon index of the Comptonized component of the spectrum, $\Gamma \approx 2$. We also suggest that *the index stability effect is an intrinsic signature of atolls, for example GX 3+1, 4U 1728-34, whereas in BHs Γ monotonically increases with $\dot{M}$ and finally saturates at high $\dot{M}$ values* (see ST09). The indices of BHCs (GRO J1655-40, GX 339-4, GRS 1915+105, H1743-322, SS 433, Cyg X-1, 4U 1543-47, XTE J1550-564) demonstrate strong correlation with $\dot{M}$ or with the soft (disk) photon normalization $N_{com} = L_{39}/D_{10}^2 \propto \dot{M}$. When $\dot{M}$ exceeds a certain level (a few times

of the Eddington mass accretion rate) the index Γ starts to saturate. In this Paper we show that the behavior of the index Γ for NS GX 340+0 is the same to that in *atolls*, 4U 1728-34, GX 3+1, and thus is drastically different from that in BHCs.

Our observational results and their theoretical explanation supports the idea that the X-ray spectral evolution of NS LMXBs, reveal the unique quasi-stability of the photon indices Γ of each Comptonized components near a value of 2. This can be explained by the model in which the energy release in the transition layer is much higher than that in the accretion disk and consequently the NS blackbody and thermal Comptonized components are dominant in the X-ray emergent spectrum.

We acknowledge productive discussion of the paper with Chris Shrader. Also we should point out a very deep analysis of our paper by the referee who suggests many important modifications of the paper.

REFERENCES

- Alpar M. A., Hasinger G., Shaham J., Yancopoulos S., 1992, A&A 257, 627
- Bisnovatyi, G. S., Khlopov, M.Yu., Chechetkin, V.M. & Eramzhyan 1980, Sov. Astr., 24, 716
- Boella G., Butler R. C., Perola G. C., et al., 1997, A&A, 122, 299
- Bradt, H. V., Rothschild, R. E. & Swank, J. H. 1993, A&AS, 97, 355
- Church, M. J., Halai, G. S. & Bałucińska-Church, M. 2006, A&A, 460, 233
- Corbet R. H. D., Smale A. P., Charles P. A., Lewin W. H. G., Menzies J. W., Naylor T., Penninx, W., Sztajno M., Thorstensen J. R., Trümper J., van Paradijs J., 1989, MNRAS 239, 533
- Cowley, A. P., Crampton, D., Hutchings, J. B., 1979, ApJ, 231, 539
- D’Aí, A., Iaria, R., Di Salvo, T., Matt, G. and Robba, N. R. 2009 ApJ, 693, L1
- Di Salvo, T., Mendez, M., van der Klis, M., Ford, E. & Robba, N.R. 2001 ApJ, 546, 1107
- Dieters S., van der Klis M., 2000, MNRAS, 311, 201
- Farinelli, R., Titarchuk, L., Paizis, A. & Frontera, F. 2008, ApJ, 680, 602 (F08)
- Farinelli, R., Paizis, A. & Landi, R. and Titarchuk, L., 2009, A&A, 498, 509

- Farinelli, R. & Titarchuk, L., 2011, A&A, 525, 102 (FT11)
- Fender, R. P., & Hendry, M. A., 2000, MNRAS, 317, 1
- Ford, E. C., van der Klis, M., Méndez, M., Wijnands, R., Homan, J., Jonker, P. G., & van Paradijs, J. 2000, ApJ, 537, 368
- Ford, E.C., van der Klis, M. 1998, ApJ, 506, L39
- Ford, E. C., van der Klis, M., Kaaret, P. 1998, ApJ, 498, 41
- Fortner B., Lamb F. K., Miller G. S., 1989, Nature 342, 775
- Friedman, H., Byram E. T., and Chubb, T. A., 1967, Science, 156, 374
- Frontera, F. et al. 1997, A&AS, 122, 357
- Hasinger, G., van der Klis M., Ebisawa K., Dotani T., Mitsuda K., 1990, A&A 235,131
- Hasinger, G., & van der Klis, M. 1989, A&A, 225, 79
- Homan, J., et al. 2007, ApJ, 656, 420
- Homan, J., van der Klis, M., Wijnands, R., Vaughan, B., & Kuulkers, E. 1998, ApJ, 499, 41
- Iaria, R., et al. 2006, Chin. J. Astron. Astrophys., Suppl. 1, 6, 257
- Jonker, P. G., Wijnands R., van der Klis, M., Psaltis, D., Kuulkers, E. & Lamb, F.K. 1998, ApJ, 499, L191
- Jonker, P. G., van der Klis, M., Wijnands R., et al. 2000, ApJ, 537, 374
- Kuulkers, E & van der Klis, M. 2000, A&A, 356, L45
- Kuulkers, E., van der Klis, M., Oosterbroek, T., Asai, K., Dotani, T., van Paradijs, J. & Lewin, W. H. G., 1994, A&A, 289, 795
- Kuulkers, E. & van der Klis, M., 1996, A&A, 314, 567
- Lamb F. K., 1989, in: Two-Topics in X-ray, 23rd ESALAB Symp., J. Hunt, & B. Battrick (eds.), 1989, ESA SP-296, p. 21
- Laurent, P. & Titarchuk, L. 2007, ApJ, 656, 1056
- Lavagetto, G., et al. 2004, Nuclear Physics B Proc. Supplements, 132, 616

- Lin, D., Remillard, R.A. & Homan, J. 2009, ApJ, 696, 1257
- Margon B. et al., 1971, ApJL, 169, L45
- Méndez, M., van der Klis, M., Ford, E. C., Wijnands, R., & van Paradijs, J. 1999, ApJ, 511, L49
- Middleditch J., Priedhorsky W. C, 1986, ApJ 306, 230
- Miller, B. W., Margon, B., Burton, M. G., 1993, AJ, 106, 404
- Muno, M. P., Remillard, R. A. & Chacrabarty, D., 2002, ApJ, 568, L35
- Narayan, R & Heyl, J.S. 2002, ApJ, 574, L139
- Oosterbroek, T., van der Klis M., Kuulkers, E., van Paradijs, M., & Lewin, W.H.G. 1995, A&A, 297, 141
- Oosterbroek, T., Lewin, W. H. G., van Paradijs, M., van der Klis M., Penninx, W. & Dotani, T. 1994, A&A, 281, 803
- Parmar, A. N., et al. 1997, A&AS, 122, 309
- Penninx, W., Zwarthoed, G. A. A., van Paradijs, J., et al., 1993, A&A, 267, 92
- Penninx, W., Lewin, W.H.G., Tan, J., Mitsuda, K., van der Klis, M., & van Paradijs, J. 1991, MNRAS., 249, 113
- Penninx, W., Lewin W. H. G., Mitsuda K., van der Klis M., Van Paradijs J., Zijlstra A. A., 1990, MNRAS 243, 114
- Ponman, T., 1982, MNRAS, 201, 769
- Psaltis, D., Belloni, T., van der Klis, M. 1999, ApJ, 520, 262
- Rappaport, S., Zaumen, W., Doxsey, R., Mayer, W., 1971, ApJ, 169, L93
- Revnivtsev, M., Borozdin, K., Emelyanov, A. 1999, A&A,
- Schulz, N. S., Hasinger, G., Truemper, J., 1989, A&A, 225, 48
- Schulz, N. S., Wijers, R. A. M. J., 1993, A&A, 273, 123 (SW93)
- Seifina, E., Titarchuk, L. & Frontera, F. 2013, ApJ, 766, (STF13)
- Seifina, E. & Titarchuk, L. 2012, ApJ, 747, 99 (ST12)

- Seifina, E. & Titarchuk, L. 2011, ApJ, 738, 128 (ST11)
- Seifina, E. & Titarchuk, L. 2010, ApJ, 722, 586 (ST10)
- Shakura, N.I., & Sunyaev, R.A. 1973, A&A, 24, 337
- Shaposhnikov, N. & Titarchuk, L. 2009, ApJ, 699, 443 (ST09)
- Shirey, R. E., Bradt, H. V. & Levine, A. M., 1999, ApJ, 517, 472
- Strohmayer, T. 1998, in American Institute of Physics Conference Series, 397 (astro-ph/9802022v1)
- Strohmayer, T. E., Zhang, W., Swank, J. H., Smale, A., Titarchuk, L., & Day, C. 1996, ApJ, 469, L9
- Titarchuk L., 1994, ApJ, 434, 570
- Titarchuk, L., Kuznetsov, S.I. & Shaposhnikov, N. 2007, ApJ, 667, 404
- Titarchuk, L., Lapidus, I.I. & Muslimov, A. 1998, ApJ, 419, 315
- Titarchuk, L., & Lyubarskij, Y. 1995, ApJ, 450, 876
- Titarchuk, L. & Osherovich, V. 2001, ApJ, 555, L55
- Titarchuk, L.G. & Osherovich, V.A. 1999, ApJ, 518, L95
- Titarchuk, L. & Seifina, E. 2009, ApJ, 706, 1463
- Titarchuk, L. & Shaposhnikov, N. 2010, ApJ, 724, 1147 (TS10)
- Ueda, Y., Mitsuda, K., Murakami, H., & Matsushita, K. 2005, ApJ, 620, 274
- van der Klis, M. 2000, ARA&A, 38, 717
- van der Klis, M., Wijnands, R.A.D., Horne, K., & Chen, W. 1997, ApJ, 481, L97
- van der Klis, M., Swank, J. H., Zang, W., Jahoda, K, Morgan, E. H., Lewin, W H. G., Vaughan, B., & van Paradijs, 1996, ApJ, 469, L1
- van der Klis, M., 1994, A&A, 283, 46
- van der Klis, M., 1994, ApJS, 92, 511
- van der Klis, M., 1989, ARA&A 27, 51

- van der Klis M., Jansen F., Van Paradijs J., Lewin W. H. G., Sztjano M., Tr’umper J., 1987, ApJ, 313, L19
- van Paradijs, J., Hasinger, G., Lewin, W. H. G., et al., 1988, MNRAS, 231, 379
- van Paradijs, J. 1978, Nature, 274, 650
- van Straaten, S., Ford, E., van der Klis, M., Méndez, M. & Kaaret, Ph. 2000, ApJ, 540, 1049
- Vrtilek, S. D., Raymond, J. C, Garcia, M. R., Verbunt, F., Hasinger, G., Kürster, M., 1990, A&A, 235, 162
- Wijnands, R., van der Klis, & Rijkhorst E. 1999, ApJ, 512, L39
- Wijnands, R., & van der Klis, 1999, ApJ, 512, L39
- Wijnands, R., van der Klis, M. 1998, Nature, 394, 344
- Wijnands, R. A. D. van der Klis, M., Méndez, M., van Paradijs, J., Lewin, W. H. G., Lamb, F. K., Vaughan, B. & Kuulkers, E. 1998, ApJ, 495, 39
- Wijnands, R. A. D. van der Klis, M., Kuulkers, E., Asai, K. & Hasinger, G. 1997, A&A, 323, 399
- Zang, W. et al. 1998, ApJ, 500, L167
- Zel’dovich, Ya. B. & Shakura N. I. 1969, Sov. Astr. 13, 175

Table 1. The list of *BeppoSAX* observations of GX 340+0 used in our analysis.

Number of set	Obs. ID	Start time (UT)	End time (UT)	MJD interval
S1	20261005	1997 Sep. 4 11:53:58	1997 Sep. 4 19:33:21	50695.5-50695.9
S2	20261006	1997 Oct. 2 20:32:50	1997 Oct. 3 04:16:33	50723.8-50724.4
S3	21240001	2000 Aug. 16 05:50:34	2000 Aug. 18 05:56:11	51772.2-51774.2 ¹
S4	212400011	2000 Aug. 18 06:16:17	2000 Aug. 20 14:21:16	51774.2-51776.9 ¹
S5	212400012	2001 March 9 19:27:20	2001 March 11 11:13:47	51977.8-51979.9 ¹
S6	21375002	2001 Aug. 9 05:01:20	2001 Aug. 10 08:56:49	52130.2-52131.9 ²
Reference. (1) Iaria et al. (2006); (2) Lavagetto et al. (2004)				

Table 2. The list of groups of *RXTE* observation of GX 340+0

Number of set	Dates, MJD	RXTE Proposal ID	Dates UT	Rem.
R1	50555	20054	Apr. 17 13:40:48 – 13:08:00, 1997	
R2	50605-50609	20059	June 6 – 10, 1997	
R3	50712-50756	20053 ¹	Sept. 21 – Nov. 4, 1997	<i>BeppoSAX</i>
R4	51130-51131	30040	Nov. 13 – 14, 1998	
R5	51772-51776	50016	Aug. 16 – 20, 2000	
R6	51920-51924	50016	Jan. 11 – 15, 2001	
R7	52908-52915	80020	Sept. 26 – Oct. 3, 2003	
R8	53210-53262	80020	July 24 – Sept. 14, 2004	
R9	53874-53907	91152	May 19 – June 21, 2006	
R10	54297	93046	July 16 01:14:56 – 01:29:20, 2007	
R11	54893-54894	94312	March 3 – 4, 2009	
References: (1) Church et al. (2006)				

Table 3. Best-fit parameters of spectral analysis of BeppoSAX observation of GX 340+0 in 0.3 – 150 keV energy range in three additive models[†]: $wabs^*(Blackbody + Comptb + Gaussian)$, $wabs^*(Blackbody1+Blackbody2+Comptb+Gaussian)$ and $wabs^*(Blackbody + Comptb1 + Comptb2 + Gaussian)$.

Model	Parameter	00-20261005	00-20261006	00-21240001	00-212400011	00-212400012	00-21375002
wabs	N_H (cm^{-2})	6.38±0.09	6.18±0.16	6.52±0.13	6.56±0.09	6.53±0.09	6.29±0.09
bbody	kT_{BB} (keV)	0.99±0.02	0.98±0.03	0.69±0.02	0.98±0.03	0.68±0.01	0.60±0.03
	$N_{BB}^{\dagger\dagger}$	7.67±0.01	7.41±0.06	10.43±0.07	7.56±0.02	7.41±0.05	10.41±0.07
comptb	$\alpha = \Gamma - 1$	1.02±0.01	1.00±0.05	1.09±0.04	1.02±0.03	1.03±0.01	1.08±0.06
	kT_s (keV)	1.56±0.04	1.49±0.05	1.76±0.06	0.48±0.04	1.43±0.04	1.40±0.04
	$\log A$	-1.29±0.03	-0.86±0.02	-0.98±0.03	-0.86±0.03	-0.72±0.07	-0.86±0.05
	kT_e (keV)	6.21±0.07	3.98±0.08	4.89±0.08	3.62±0.06	3.98±0.08	4.98±0.09
	$N_{Com}^{\dagger\dagger}$	7.47±0.05	10.02±0.01	15.19±0.09	10.02±0.07	14.20±0.04	13.61±0.05
Gaussian	E_{line} (keV)	6.87±0.06	6.80±0.08	6.74±0.07	7.00±0.06	6.80±0.06	6.71±0.05
	σ_{line} (keV)	0.65±0.05	0.60±0.01	0.81±0.06	0.69±0.04	0.91±0.06	0.74±0.05
	$N_{line}^{\dagger\dagger}$	0.45±0.02	0.43±0.06	0.55±0.07	0.26±0.07	0.39±0.07	0.35±0.08
	χ^2_{red} (d.o.f.)	3.2 (336)	1.29 (330)	1.59 (332)	1.3 (332)	1.43(332)	2.7 (330)
wabs	N_H (cm^{-2})	5.40±0.07	6.29±0.06	6.43±0.09	6.43±0.09	6.35±0.07	5.70±0.09
bbody1	kT_{BB1} (keV)	0.60±0.08	0.59±0.04	0.61±0.05	0.52±0.06	0.58±0.08	0.60±0.02
	$N_{BB1}^{\dagger\dagger}$	2.1±0.1	7.58±0.03	7.98±0.09	7.81±0.06	8.2±0.1	3.70±0.09
bbody2	kT_{BB2} (keV)	1.10±0.03	1.50±0.06	1.51±0.02	1.47±0.04	1.51±0.02	1.50±0.04
	$N_{BB2}^{\dagger\dagger}$	2.00±0.07	6.42±0.5	9.95±0.08	7.96±0.09	6.49±0.08	4.75±0.09
comptb	$\alpha = \Gamma - 1$	1.00±0.01	1.06±0.04	0.99±0.03	1.02±0.04	1.05±0.06	1.10±0.09
	kT_s (keV)	1.21±0.09	1.20±0.02	1.26±0.07	0.18±0.06	1.20±0.08	1.19±0.08
	$\log A$	-1.02±0.06	-0.55±0.02	-0.21±0.03	-0.29±0.02	-1.08±0.06	2.0 ^{†††}
	kT_e (keV)	4.66±0.08	3.51±0.09	3.54±0.04	3.35±0.03	4.33±0.09	3.32±0.02
	$N_{Com}^{\dagger\dagger}$	10.78±0.09	8.67±0.05	6.24±0.09	7.16±0.08	7.63±0.06	5.43±0.07
Gaussian	E_{line} (keV)	6.75±0.08	6.78±0.09	6.80±0.07	6.81±0.05	6.90±0.07	6.75±0.05
	σ_{line} (keV)	0.95±0.04	0.90±0.06	0.89±0.07	0.79±0.04	0.91±0.05	0.94±0.03
	$N_{line}^{\dagger\dagger}$	2.60±0.05	0.66±0.09	0.65±0.08	0.63±0.06	0.53±0.07	0.65±0.09
	χ^2_{red} (d.o.f.)	3.28 (334)	1.07 (328)	1.53 (330)	1.65 (330)	1.63(330)	2.18 (327)
wabs	N_H (cm^{-2})	5.7±0.1	6.0±0.1	6.01±0.03	6.3±0.1	6.4±0.1	5.4±0.3
bbody	kT_{BB} (keV)	0.56±0.08	0.57±0.04	0.56±0.03	0.53±0.05	0.43±0.04	0.57±0.01
	$N_{BB}^{\dagger\dagger}$	0.92±0.09	0.9±0.1	0.90±0.05	0.91±0.04	4.47±0.02	2.01±0.06
comptb1	$\alpha_1 = \Gamma_1 - 1$	0.99±0.04	1.00±0.08	1.02±0.07	1.01±0.07	0.99±0.01	1.04±0.06
	kT_{s1} (keV)	0.70±0.03	0.6±0.1	0.69±0.02	0.70±0.03	0.83±0.04	1.03±0.01
	$\log A_1$	-0.32±0.06	-0.30±0.07	-0.30±0.05	-0.30±0.03	-1.42±0.06	-1.33±0.07
	$kT_e^{(1)}$ (keV)	5.38±0.02	5.42±0.03	5.60±0.04	5.41±0.04	6.04±0.03	19.50±0.07
	$N_{Com1}^{\dagger\dagger}$	1.36±0.07	1.86±0.04	0.57±0.06	1.53±0.04	9.96±0.06	3.97±0.07
comptb2	$\alpha_2 = \Gamma_2 - 1$	1.01±0.03	1.04±0.05	1.01±0.03	1.05±0.08	1.06±0.05	0.99±0.02
	kT_{s2} (keV)	1.0±0.1	1.00±0.09	1.71±0.05	1.06±0.09	1.50±0.02	1.51±0.04
	$kT_e^{(2)}$ (keV)	1.71±0.09	1.90±0.02	2.04±0.08	2.05±0.03	1.52±0.03	2.76±0.09
	$N_{Com2}^{\dagger\dagger}$	10.84±0.07	12.17±0.06	14.26±0.05	13.11±0.04	11.79±0.05	8.20±0.06

Table 3—Continued

Model	Parameter	00-20261005	00-20261006	00-21240001	00-212400011	00-212400012	00-21375002
Gaussian	E_{line} (keV)	6.85 ± 0.08	6.68 ± 0.03	6.68 ± 0.05	6.70 ± 0.04	6.71 ± 0.03	6.80 ± 0.06
	$N_{line}^{\dagger\dagger}$	1.07 ± 0.09	0.89 ± 0.08	0.68 ± 0.09	0.77 ± 0.09	0.89 ± 0.02	0.7 ± 0.1
	χ_{red}^2 (d.o.f.)	0.97 (332)	0.98 (326)	0.99 (328)	1.38 (328)	1.49 (328)	1.28 (326)

[†] Errors are given at the 90% confidence level. ^{††} The normalization parameters of *Blackbody* and *CompTB* components are in units of L_{37}^{soft}/d_{10}^2 erg/s/kpc², where L_{37}^{soft} is the soft photon luminosity in units of 10^{37} erg/s, d_{10} is the distance to the source in units of 10 kpc and *Gaussian* component is in units of $10^{-2} \times total \text{ photons cm}^{-2} \text{ s}^{-1}$ in line; ^{†††} when parameter $\log(A) \gg 1$, it is fixed to a value 2.0 for the model $wabs * (Blackbody1 + Blackbody2 + Comptb + Gaussian)$, parameter $\log(A_2)$ is fixed at 2.0 for the model $wabs * (Blackbody + Comptb1 + Comptb2 + Gaussian)$ (see comments in the text); σ_{line} of *Gaussian* component is fixed to a value 0.60 keV for the model $wabs * (Blackbody + Comptb1 + Comptb2 + Gaussian)$ (see comments in the text), N_H is units of 10^{22} cm^{-2} .

Table 4. Best-fit parameters of spectral analysis of PCA+HEXTE/*RXTE* observations of GX 340+1 in 3 – 150 keV energy range[†]. Adopted model: *wabs*(Blackbody + Comptb1 + Comptb2 + Gaussian)*. Parameter errors are given at 90% confidence level.

Observational ID	MJD, day	$\alpha_1 =$ $\Gamma_1 - 1$	$kT_e^{(1)}$, keV	$\log(A_1)$	$N_{Com1}^{\dagger\dagger\dagger}$	$N_{Bbody}^{\dagger\dagger\dagger}$	$\alpha_2 =$ $\Gamma_2 - 1$	$kT_e^{(2)}$, keV	$N_{Com2}^{\dagger\dagger\dagger}$	E_{line} , keV	$N_{line}^{\dagger\dagger\dagger}$	χ_{red}^2 (d.o.f.)	$F_1/F_2^{\dagger\dagger\dagger}$
20054-04-01-000	50555.570	0.99(1)	17.96(5)	-1.19(4)	4.72(9)	1.03(9)	1.01(1)	2.72(7)	11.77(3)	6.74(9)	1.05(2)	1.17(67)	9.89/3.49
20059-01-01-000	50605.255	0.99(2)	5.43(2)	-0.54(2)	2.06(6)	0.92(3)	1.02(2)	1.63(4)	13.29(5)	6.45(8)	2.04(3)	1.46(67)	9.07/1.14
20059-01-01-01	50605.904	1.03(3)	18.77(5)	-1.32(2)	4.68(4)	1.04(8)	0.99(2)	2.74(1)	11.72(6)	6.67(9)	1.03(5)	0.93(67)	9.87/3.39
20059-01-01-02	50606.471	1.03(2)	20.40(9)	-0.90(3)	3.14(9)	1.06(5)	1.00(1)	2.91(6)	13.4(9)	6.50(7)	1.02(4)	1.08(67)	11.61/3.59
20059-01-01-03	50606.996	1.04(3)	12.32(8)	-1.09(4)	3.45(5)	1.01(6)	1.03(4)	2.49(1)	16.03(4)	6.5(1)	2.09(2)	1.19(67)	12.36/3.50
20059-01-01-04	50607.329	1.02(2)	12.34(9)	-0.06(3)	3.50(6)	1.07(4)	1.00(2)	2.48(4)	16.45(9)	6.4(1)	2.03(2)	1.31(67)	12.61/3.55
20059-01-01-05	50608.008	0.99(1)	15.08(5)	-1.29(3)	6.47(5)	1.09(5)	1.02(3)	2.70(2)	11.65(9)	6.40(9)	2.83(6)	1.08(67)	10.86/3.53
20059-01-01-06	50609.062	1.01(2)	11.28(8)	-1.05(1)	4.03(6)	1.08(6)	1.01(1)	2.27(1)	17.03(9)	6.5(1)	2.29(8)	0.94(67)	11.70/2.71
20053-05-01-00	50712.048	1.02(3)	4.01(9)	-0.42(5)	2.88(9)	0.92(7)	0.98(3)	2.18(3)	8.67(8)	6.74(3)	0.75(7)	0.84(67)	10.58/1.56
20053-05-01-01	50714.179	1.06(4)	3.12(4)	-0.11(2)	2.05(6)	0.92(2)	1.01(1)	1.85(2)	10.1(1)	6.41(8)	2.31(6)	0.97(67)	10.00/1.88
20053-05-01-02	50716.066	1.02(2)	16.65(4)	-1.20(3)	4.76(5)	1.03(9)	1.02(3)	2.69(1)	12.55(4)	6.7(1)	1.03(9)	1.09(67)	10.52/3.47
20053-05-01-03	50920.197	1.05(3)	12.96(9)	-1.08(4)	3.62(9)	1.01(8)	1.01(5)	2.59(7)	13.11(6)	6.4(1)	2.1(1)	0.94(67)	10.51/3.12
20053-05-02-00	50753.947	0.99(1)	4.43(5)	-0.56(1)	2.04(2)	0.92(1)	1.04(3)	1.56(8)	16.35(1)	6.62(2)	0.70(4)	1.12(67)	9.42/1.14
20053-05-02-01	50754.091	0.99(2)	10.09(8)	-1.14(3)	5.31(3)	1.07(5)	1.02(2)	2.26(2)	12.80(9)	6.57(9)	2.28(5)	1.07(67)	10.82/2.29
20053-05-02-02	50754.825	1.02(2)	12.30(9)	-1.08(4)	3.38(6)	1.0(1)	1.01(3)	2.48(1)	15.98(8)	6.36(9)	2.09(9)	0.93(67)	12.37/3.39
20053-05-02-03	50755.133	1.03(2)	11.34(8)	-1.09(3)	3.41(5)	1.07(5)	1.03(4)	2.45(2)	15.54(7)	6.47(8)	2.08(4)	0.89(67)	12.04/3.19
20053-05-02-05	50756.680	1.01(1)	17.85(4)	-1.21(2)	4.76(7)	1.04(8)	1.03(5)	2.71(2)	11.71(3)	6.8(1)	1.06(5)	0.89(67)	9.89/3.28
20053-05-02-04	50756.685	1.05(4)	12.94(5)	-1.08(4)	3.40(8)	1.0(1)	0.99(6)	2.54(1)	15.81(4)	6.42(9)	2.04(9)	0.91(67)	12.23/3.57
30040-04-01-00	51130.995	1.04(2)	16.87(4)	-1.46(3)	6.23(7)	1.04(6)	1.00(2)	2.69(3)	12.03(8)	6.45(2)	2.18(2)	0.86(67)	10.19/3.39
30040-04-01-06	51131.435	1.02(2)	17.78(9)	-0.99(6)	4.79(6)	1.09(8)	1.02(3)	2.71(2)	11.31(5)	6.70(9)	1.03(4)	1.02(67)	9.70/3.48
30040-04-01-01	51131.601	1.02(3)	18.67(8)	-1.34(2)	4.70(8)	1.08(7)	1.02(3)	2.78(3)	10.71(4)	6.70(8)	1.03(2)	1.06(67)	9.18/3.24
30040-04-01-020	51131.877	1.00(1)	19.65(7)	-1.24(4)	5.01(6)	1.09(8)	1.01(1)	2.70(1)	11.83(3)	6.70(9)	1.05(3)	0.96(67)	9.99/3.38
30040-04-01-03	51132.596	1.02(2)	12.73(2)	-1.05(2)	3.48(3)	1.03(7)	1.01(2)	2.39(2)	16.86(4)	6.47(6)	2.24(2)	0.96(67)	12.91/3.32
30040-04-01-05	51132.829	1.05(3)	12.91(8)	-1.08(4)	3.45(9)	1.04(9)	1.03(3)	2.50(1)	16.03(5)	6.40(1)	2.08(1)	1.00(67)	13.27/3.72
30040-04-01-04	51132.881	1.01(1)	12.93(1)	-1.07(1)	3.33(5)	1.03(8)	1.02(4)	2.51(2)	16.68(9)	6.40(6)	2.14(2)	0.96(67)	12.67/3.49
50016-01-01-11	51772.077	0.99(2)	12.91(4)	-1.08(4)	3.40(7)	1.06(3)	0.99(3)	2.49(1)	16.51(5)	6.4(1)	2.95(3)	0.71(67)	13.24/3.96
50016-01-01-12	51772.144	1.04(4)	11.8(1)	-1.10(7)	3.5(1)	1.01(4)	1.01(2)	2.4(1)	17.29(9)	6.41(7)	2.18(4)	1.03(67)	13.11/3.42

Table 4—Continued

Observational ID	MJD, day	$\alpha_1 =$ $\Gamma_1 - 1$	$kT_e^{(1)}$, keV	$\log(A_1)$	$N_{Com1}^{\dagger\dagger\dagger}$	$N_{Bbody}^{\dagger\dagger\dagger}$	$\alpha_2 =$ $\Gamma_2 - 1$	$kT_e^{(2)}$, keV	$N_{Com2}^{\dagger\dagger\dagger}$	E_{line} , keV	$N_{line}^{\dagger\dagger\dagger}$	χ_{red}^2 (d.o.f.)	$F_1/F_2^{\dagger\dagger\dagger}$
50016-01-01-000	51772.210	0.99(2)	12.72(9)	-1.04(2)	3.39(9)	1.04(9)	1.03(4)	2.45(3)	16.55(8)	6.70(9)	0.80(5)	1.02(67)	12.69/3.43
50016-01-01-01	51772.708	1.01(2)	12.28(8)	-1.09(2)	3.49(5)	1.04(8)	0.98(4)	2.39(1)	17.34(5)	6.4(1)	2.62(6)	1.12(67)	13.08/3.52
50016-01-01-10	51772.742	1.01(1)	12.96(2)	-1.08(4)	3.35(7)	1.02(3)	0.99(1)	2.54(2)	16.19(8)	6.42(9)	2.89(2)	1.01(67)	12.50/3.73
50016-01-01-07	51773.012	1.02(2)	11.06(9)	-1.06(7)	4.0(1)	1.01(6)	1.01(2)	2.29(1)	19.4(1)	6.53(8)	2.29(5)	0.76(67)	12.83/3.10
50016-01-01-02	51773.082	1.04(3)	12.9(1)	-1.08(6)	3.48(9)	1.04(9)	1.01(1)	2.45(5)	17.31(6)	6.40(7)	2.16(7)	0.76(67)	13.11/3.75
50016-01-01-03	51773.146	1.02(2)	12.3(1)	-1.02(3)	3.39(4)	1.32(5)	1.05(4)	2.45(9)	17.46(5)	6.70(8)	2.94(5)	0.87(67)	12.72/3.81
50016-01-01-04	51773.211	1.01(1)	12.26(9)	-1.10(2)	3.47(5)	1.33(9)	1.04(3)	2.46(7)	17.67(8)	6.64(9)	2.90(4)	0.76(67)	12.76/3.77
50016-01-01-05	51773.281	0.99(2)	13.0(1)	-1.06(2)	4.49(3)	1.02(8)	1.05(4)	2.46(1)	17.41(9)	6.40(5)	2.79(5)	1.16(67)	12.61/3.71
50016-01-01-09	51773.362	1.02(3)	11.09(3)	-1.13(1)	5.6(1)	1.12(6)	1.01(1)	2.32(2)	14.31(9)	6.54(9)	2.09(2)	1.07(67)	11.47/2.69
50016-01-01-06	51773.619	0.99(1)	10.0(2)	-1.14(9)	5.5(1)	1.05(2)	1.00(3)	2.22(3)	12.51(3)	6.6(1)	2.15(3)	1.06(67)	10.65/2.23
50016-01-01-08	51773.939	1.05(4)	9.2(1)	-1.13(4)	5.4(1)	1.1(1)	1.03(4)	2.13(9)	13.3(1)	6.5(1)	2.39(9)	1.24(67)	10.82/2.09
50016-01-02-00	51774.005	1.02(2)	5.42(4)	-0.99(3)	1.39(8)	0.92(4)	0.99(2)	1.52(2)	15.5(1)	6.87(9)	0.56(3)	1.39(67)	9.25/0.98
50016-01-02-18	51774.075	1.05(6)	3.30(5)	-0.32(2)	1.31(9)	0.92(9)	0.99(1)	1.52(3)	15.92(8)	6.89(4)	0.56(3)	1.28(67)	9.04/0.99
50016-01-02-05	51774.138	1.01(2)	2.8(1)	-1.08(2)	3.5(1)	1.1(1)	0.99(3)	1.55(1)	13.05(7)	6.39(5)	1.90(4)	1.26(67)	9.56/1.04
50016-01-02-03	51774.204	0.99(1)	3.67(6)	-0.97(6)	1.32(8)	0.93(8)	1.02(2)	1.51(2)	17.5(1)	6.89(4)	0.59(5)	1.27(67)	9.64/1.04
50016-01-02-16	51774.270	1.01(3)	3.34(3)	-0.11(3)	1.39(7)	0.91(4)	1.01(1)	1.70(8)	15.60(9)	6.67(5)	0.57(9)	1.22(67)	8.89/1.32
50016-01-02-04	51774.574	1.01(2)	12.9(1)	-1.08(2)	3.5(1)	1.1(1)	0.99(4)	2.46(1)	15.23(6)	6.40(6)	2.86(4)	0.99(67)	12.38/3.51
50016-01-02-12	51774.704	0.99(1)	11.04(3)	-1.05(4)	4.08(9)	1.01(7)	1.01(3)	2.36(2)	16.94(5)	6.33(5)	4.28(5)	0.95(67)	11.47/2.87
50016-01-02-17	51775.001	1.02(2)	11.07(9)	-1.13(5)	5.49(7)	1.06(3)	1.00(1)	2.4(1)	14.82(6)	6.58(6)	2.94(8)	0.98(67)	11.88/2.94
50016-01-02-02	51775.067	1.07(6)	11.1(1)	-1.08(2)	4.02(8)	1.9(1)	1.01(2)	2.4(1)	16.41(8)	6.38(7)	3.10(1)	0.97(67)	12.54/3.18
50016-01-02-07	51775.200	1.02(2)	11.08(9)	-1.05(4)	4.18(9)	1.07(9)	1.00(3)	2.3(1)	16.53(8)	6.37(4)	2.02(3)	1.03(67)	12.02/3.05
50016-01-02-06	51775.267	1.04(3)	11.1(1)	-1.12(3)	4.87(2)	1.00(1)	1.02(2)	2.39(9)	14.45(6)	6.56(3)	2.08(2)	1.02(67)	11.17/2.62
50016-01-02-15	51775.533	1.00(1)	12.9(1)	-1.09(1)	3.49(9)	1.02(3)	1.05(5)	2.45(6)	15.56(3)	6.42(7)	2.06(3)	0.97(67)	11.99/3.32
50016-01-02-08	51775.734	1.05(4)	12.87(8)	-1.08(5)	3.41(5)	1.03(2)	0.99(3)	2.49(1)	15.52(8)	6.45(4)	2.88(4)	0.99(67)	12.02/3.40
50016-01-02-10	51775.998	1.00(1)	12.9(1)	-1.07(4)	3.52(9)	1.07(5)	1.06(4)	2.39(2)	15.15(8)	6.40(5)	2.49(5)	0.94(67)	12.31/3.39
50016-01-02-09	51776.064	1.01(2)	12.84(9)	-1.06(5)	3.34(7)	1.0(1)	0.99(3)	2.41(1)	17.31(9)	6.40(6)	2.69(3)	0.99(67)	12.42/3.37
50016-01-02-13	51776.131	1.02(3)	11.07(8)	-1.13(7)	5.58(9)	1.06(4)	1.01(5)	2.31(3)	14.21(9)	6.6(1)	2.75(4)	1.20(67)	11.53/2.72

Table 4—Continued

Observational ID	MJD, day	$\alpha_1 =$ $\Gamma_1 - 1$	$kT_e^{(1)},$ keV	$\log(A_1)$	$N_{Com1}^{\dagger\dagger\dagger}$	$N_{Bbody}^{\dagger\dagger\dagger}$	$\alpha_2 =$ $\Gamma_2 - 1$	$kT_e^{(2)},$ keV	$N_{Com2}^{\dagger\dagger\dagger}$	$E_{line},$ keV	$N_{line}^{\dagger\dagger\dagger}$	χ_{red}^2 (d.o.f.)	$F_1/F_2^{\dagger\dagger\dagger}$
50016-01-02-11	51776.529	0.99(2)	3.24(2)	-0.38(6)	2.07(9)	0.92(6)	1.03(4)	1.94(8)	10.23(7)	6.43(7)	2.31(9)	0.92(67)	9.10/1.59
50016-01-02-01	51776.680	1.01(2)	4.12(4)	-0.46(3)	2.01(7)	0.9(1)	1.02(3)	1.8(1)	11.1(1)	6.35(6)	2.26(3)	1.16(67)	8.85/1.34
50016-01-04-01G	51920.397	1.01(1)	18.81(9)	-1.81(5)	4.78(8)	1.02(4)	1.01(2)	2.9(1)	11.24(9)	6.7(1)	1.08(5)	0.87(67)	9.28/3.61
50016-01-04-00	51920.705	1.03(4)	15.92(5)	-1.64(3)	6.26(9)	1.08(9)	0.99(1)	2.69(9)	13.71(9)	6.41(4)	2.10(3)	1.03(67)	11.64/3.78
50016-01-03-11	51921.001	0.99(2)	12.94(7)	-1.07(9)	3.47(9)	1.03(4)	1.01(2)	2.49(1)	16.27(5)	6.46(5)	2.14(4)	1.12(67)	12.47/3.61
50016-01-03-12	51921.317	0.99(3)	11.79(4)	-1.08(5)	5.12(8)	1.06(1)	1.00(1)	2.33(4)	15.54(4)	6.53(5)	2.29(3)	0.69(67)	12.23/2.98
50016-01-04-02	51921.384	1.02(2)	4.35(2)	-0.45(4)	2.05(7)	0.93(3)	1.01(2)	1.65(5)	13.2(1)	6.50(2)	2.08(5)	0.97(67)	8.92/1.24
50016-01-03-02	51921.700	0.98(2)	3.44(3)	-0.36(2)	2.19(9)	0.92(4)	0.98(2)	1.68(1)	14.09(8)	6.49(5)	2.11(2)	1.12(67)	9.79/1.41
50016-01-03-01	51922.039	1.01(1)	3.08(9)	-0.07(5)	3.17(9)	1.04(5)	0.97(4)	2.42(2)	17.68(5)	6.46(4)	2.00(3)	1.18(67)	10.01/2.09
50016-01-03-13	51922.312	1.03(5)	11.3(1)	-1.14(9)	3.39(8)	1.07(6)	1.03(4)	2.62(8)	12.51(9)	6.54(7)	1.01(2)	0.79(67)	10.84/1.96
50016-01-03-00	51922.379	1.05(4)	5.42(2)	-0.69(3)	2.06(7)	0.95(5)	1.01(2)	1.58(1)	13.34(7)	6.7(1)	0.73(5)	1.48(67)	8.52/1.02
50016-01-03-14	51922.648	1.04(3)	8.40(3)	-1.32(6)	1.92(6)	0.96(2)	0.99(1)	1.55(4)	14.50(8)	6.71(7)	0.77(8)	1.28(67)	8.47/0.91
50016-01-03-03	51922.702	1.05(4)	4.04(9)	-0.41(5)	2.09(5)	0.92(3)	0.98(4)	1.84(2)	12.08(9)	6.42(5)	2.31(2)	1.05(67)	9.49/1.54
50016-01-03-040	51922.733	1.01(1)	11.80(3)	-1.14(2)	5.43(6)	1.04(7)	1.02(3)	2.29(3)	14.19(2)	6.50(5)	2.03(3)	0.83(67)	11.37/2.62
50016-01-03-05	51923.383	1.09(2)	17.67(7)	-1.55(6)	4.7(1)	1.06(5)	0.99(2)	2.80(1)	12.43(7)	6.73(4)	1.03(1)	0.90(67)	10.30/3.28
50016-01-03-06	51923.628	1.03(3)	19.91(2)	-1.02(3)	3.08(9)	3.07(3)	1.00(1)	2.74(2)	10.25(4)	6.40(5)	3.90(2)	0.97(67)	9.23/3.51
50016-01-03-070	51923.694	1.02(2)	17.26(3)	-1.51(4)	4.69(8)	1.05(3)	1.02(2)	2.76(1)	12.97(3)	6.72(6)	1.01(4)	1.02(67)	10.65/3.76
50016-01-03-09	51924.299	1.05(6)	17.10(6)	-1.34(5)	4.67(6)	1.06(2)	1.03(2)	2.66(3)	14.76(7)	6.71(5)	0.98(3)	1.16(67)	11.80/4.01
50016-01-03-10	51924.365	1.01(1)	12.52(6)	-1.07(3)	3.12(9)	1.03(4)	1.02(2)	2.44(1)	17.67(6)	6.40(5)	2.17(3)	1.00(67)	12.44/3.29
50016-01-03-08	51924.583	1.03(3)	17.51(8)	-1.59(6)	4.79(6)	0.99(20)	0.99(1)	2.8(1)	12.11(8)	6.71(7)	1.08(2)	0.79(67)	10.03/3.67
80020-02-01-00	52908.041	1.01(1)	12.92(9)	-1.07(8)	3.40(5)	1.03(9)	1.01(2)	2.44(1)	13.08(9)	6.41(2)	2.97(3)	0.98(67)	10.32/2.84
80020-02-01-01	52908.109	1.00(1)	12.25(6)	-1.05(3)	3.39(7)	1.07(8)	1.03(4)	2.42(1)	13.76(6)	6.43(2)	2.98(2)	0.94(67)	10.09/2.63
80020-02-02-00	52914.565	1.02(2)	4.24(6)	-0.53(5)	2.08(6)	0.92(5)	1.01(1)	1.67(2)	10.04(5)	6.24(9)	2.31(5)	0.89(67)	7.23/1.04
80020-02-02-01	52914.941	1.05(4)	12.93(8)	-1.08(4)	3.45(5)	1.02(8)	0.99(2)	2.48(6)	12.91(4)	6.42(9)	2.94(3)	1.07(67)	10.24/2.92
80020-02-02-02	52915.008	1.03(2)	11.02(4)	-1.13(7)	5.56(4)	1.01(7)	0.99(1)	2.31(7)	11.68(6)	6.50(9)	2.28(2)	1.02(67)	9.35/2.23
80020-02-04-00	53210.048	1.01(2)	12.94(7)	-1.08(7)	3.47(6)	1.05(6)	0.98(2)	2.53(6)	14.38(4)	6.45(3)	2.82(7)	1.33(67)	11.26/3.29
80020-02-05-00	53222.037	1.07(5)	9.79(5)	-1.12(1)	4.05(4)	1.03(7)	1.05(4)	2.12(2)	12.06(8)	6.51(9)	2.96(3)	1.06(67)	9.61/1.85

Table 4—Continued

Observational ID	MJD, day	$\alpha_1 =$ $\Gamma_1 - 1$	$kT_e^{(1)},$ keV	$\log(A_1)$	$N_{Com1}^{\dagger\dagger\dagger}$	$N_{Bbody}^{\dagger\dagger\dagger}$	$\alpha_2 =$ $\Gamma_2 - 1$	$kT_e^{(2)},$ keV	$N_{Com2}^{\dagger\dagger\dagger}$	$E_{line},$ keV	$N_{line}^{\dagger\dagger\dagger}$	χ_{red}^2 (d.o.f.)	$F_1/F_2^{\dagger\dagger\dagger}$
80020-02-06-00	53238.305	1.04(4)	18.96(8)	-1.08(5)	3.43(5)	1.04(8)	1.01(3)	2.51(4)	14.07(7)	6.40(9)	2.73(2)	1.36(67)	11.07/3.18
80020-02-07-00	53245.879	1.03(2)	10.14(9)	-1.13(2)	4.51(9)	1.08(7)	1.04(5)	2.24(2)	14.14(5)	6.53(9)	2.01(2)	1.20(67)	9.85/2.16
80020-02-08-00	53255.234	0.98(5)	20.53(2)	-0.91(3)	3.06(8)	1.07(6)	1.02(2)	3.00(1)	9.13(6)	6.39(7)	1.67(3)	1.37(67)	7.43/3.32
80020-02-09-00	53262.321	0.99(2)	11.23(8)	-1.13(4)	5.43(4)	1.04(9)	1.02(3)	2.31(1)	13.97(9)	6.51(8)	2.25(5)	1.10(67)	10.76/2.51
94312-01-01-00	54893.348	1.01(4)	17.84(8)	-1.63(9)	4.73(8)	1.0(1)	1.01(1)	2.95(2)	9.46(9)	6.70(8)	0.93(4)	1.14(67)	8.48/2.96
94312-01-01-02	54894.766	0.99(2)	14.01(9)	-1.02(5)	5.15(7)	1.06(9)	1.02(4)	2.69(3)	11.53(8)	6.46(6)	2.86(5)	1.16(67)	10.53/3.46
94312-01-01-04	54894.895	1.10(9)	12.54(8)	-1.07(3)	3.46(5)	1.1(1)	1.01(5)	2.44(1)	14.15(4)	6.40(7)	2.69(2)	1.17(67)	10.38/2.60
93046-04-01-00	54297.052	1.02(5)	11.02(9)	-1.05(2)	4.03(9)	1.02(6)	1.00(2)	2.39(1)	12.58(7)	6.3(1)	0.9(1)	1.13(67)	9.07/2.21
91152-03-01-000	53874.858	0.99(2)	9.02(8)	-1.04(6)	4.6(1)	1.05(9)	1.02(6)	2.28(4)	10.57(9)	6.58(5)	2.72(6)	1.30(67)	9.57/1.90
91152-03-02-00	53907.403	1.05(4)	10.11(1)	-1.10(5)	4.5(1)	1.07(4)	1.01(5)	2.21(1)	10.9(1)	6.53(9)	2.23(1)	1.16(67)	9.14/1.92
91152-03-02-01	53907.680	1.01(1)	11.0(1)	-1.06(9)	4.03(7)	1.06(5)	1.03(4)	2.29(2)	15.74(5)	6.43(8)	2.38(3)	0.98(67)	10.25/2.56

[†] The spectral model is $wabs * (blackbody + Comptb1 + Comptb2 + Gaussian)$; color temperature T_{BB} of *Bbody* component is fixed at 0.6 keV and “seed” photon temperatures T_{s1}/T_{s2} are fixed at 1.1/1.5 keV, respectively (see comments in the text); ^{††} parameter $\log(A_2)$ is fixed at 2.0 (see comments in the text), ^{†††} normalization parameters of *blackbody* and *COMPTB* components are in units of L_{37}/d_{10}^2 , where L_{37} is the source luminosity in units of 10^{37} erg/s, d_{10} is the distance to the source in units of 10 kpc and *Gaussian* component is in units of $10^{-2} \times total \text{ photons cm}^{-2} s^{-1}$ in line, σ_{line} of Gaussian component is fixed to a value 0.6 keV (see comments in the text), N_H was fixed at value of $6.2 \times 10^{22} \text{ cm}^{-2}$ (Iaria et al., 2006), ^{††††} spectral fluxes (F_1/F_2) in units of $\times 10^{-9} \text{ ergs/s/cm}^2$ for (3 – 10) and (10 – 50) keV energy ranges respectively.

Table 5. Comparisons of the best-fit parameters of *Z* source GX 340+0 and *atoll* sources GX 3+1¹ and 4U 1728-34²

Source name	Alternative name	Class ³	Distance, kpc	Presence of kHz QPO	kT_e , keV	N_{comptb} L_{39}^{soft}/D_{10}^2	kT_{BB} keV	kT_s keV	f
4U 1642-45	GX 340+0	Z, Sp, B	10.5 ⁴	+ ⁷	3-21	0.08-0.2	0.6	1.1-1.5	0.01-0.5
4U 1744-26	GX 3+1	Atoll, Sp, B	4.5 ⁵	none ⁸	2.3-4.5	0.04-0.15	0.6	1.16-1.7	0.2-0.9
4U 1728-34	GX 354-0	Atoll, Su, D	4.2-6.4 ⁶	+ ⁹	2.5-15	0.02-0.09	0.6-0.7	1.3	0.5-1

References: (1) ST12, (2) ST11, (3) Classification of the system in the various schemes (see text): Sp = supercritical, Su = subcritical, B = bulge, D = disk, (4) Fender & Henry (2000), Ford et al. (1998), Christian & Swank (1997), (5) Kuulkers & van der Klis (2000), Ford et al. (2000), (6) van Paradijs (1978), (7) Jonker et al. (1998), (8) Strohmayer (1998), (9) Titarchuk & Osherovich (1999)




Fig. 1.— A suggested geometry of GX 340+0. Disk and neutron star soft photons are upscattered off hotter plasma of the Transition Layer (TL) located between the accretion disk and NS surface. Some fraction of these photons is seen directly by the Earth observer. Red and blue photon trajectories correspond to soft and hard (upscattered) photons respectively. In our model two Comptonization components are considered: the first one (*Comptb1*), with “seed” (disk) photon temperature $T_{s1} \lesssim 1$ keV and the CC electron temperature $kT_e^{(1)}$ varies from 3 keV to 21 keV. The second Comptonized component (*Comptb2* with $T_{s2} \sim 1.5$ keV) and $kT_e^{(2)}$ which is related to NS surface and its inner part of the TL (boundary) layer respectively.




Fig. 2.— Photon indices Γ_{com1} and Γ_{com2} plotted vs. the normalizations of the *Comptb1* and *Comptb2* components, respectively, measured in L_{39}/D_{10}^2 units (see Table 4). *Red* and *blue* points correspond to *Comptb1* and *Comptb2* components, respectively, which are related to thermal upscattering of soft photons by plasma electrons in CC and at the NS boundary layer, respectively.




Fig. 3.— Same as in Fig. 2 but plotted vs the best-fit electron temperatures of the *Comptb1* and *Comptb2* components, respectively, measured in keV (see Table 4).




Fig. 4.— The best-fit electron temperature kT_e of *Comptb1* component (in keV), using our spectral model *wabs* * (*blackbody* + *Comptb1* + *Comptb2* + *Gaussian*), is plotted versus flux ratio $[10 - 50 \text{ keV}/3 - 10 \text{ keV}]$ during Z-state events. The horizontal dashed lines mark the branch regions of Z pattern.




Fig. 5.— *Left panel:* Flux ratio $[10-50 \text{ keV}/3-10 \text{ keV}]$ versus flux in the range of 3-50 keV for *RXTE* data. The spectral branches have been indicated by *pink* points for FB, *green* points for NB and *red* points for HB. The direction of HB→NB→FB evolution is indicated by arrow. *Right panel:* Three representative EF_E diagrams for different states along Z track of GX 340+0. Data are taken from *RXTE* observations 20053-05-01-01 (*green*, NB) 20053-05-01-02 (*violet*, FB) and 20053-05-01-00 (*red*, HB).




Fig. 6.— Spectral hardness (10 – 50 keV/3 – 50 keV) vs flux in 3 – 50 keV range of Z source GX 340+0 (*red*), *atoll* sources 4U 1728-34 (*blue*, taken from ST11) and GX 3+1 (*green*, taken from ST12), and BHC SS 433 (*violet*, taken from ST10) for *RXTE* data.




Fig. 7.— The normalizations N_{com1} (red) and N_{com2} (blue) of *Comptb1* and *Comptb2* components, respectively, are plotted versus flux ratio [10 – 50 keV/3 – 10 keV] using our spectral model $wabs*(blackbody+Comptb1+Comptb2+Gaussian)$ during evolution events.




Fig. 8.— Electron temperature $kT_e^{(1)}$ (in keV) plotted versus illumination fraction f for atoll sources GX 3+1 and 4U 1728-34 as well as for Z- source GX 340+0 (using a variable f_1 factor) during different states. Red, green and blue points correspond to *RXTE* observations of GX 340+0, GX 3+1 and 4U 1728-34 respectively. Along the track of Z source GX 340+0 we show a sequence of CCD states: HB, NB and FB. Whereas along the tracks of atoll sources 4U 1728-34 and GX 3+1 we indicate a sequence of CCD states for these particular sources. kT_e is related to the sequence of CCD states for Z-source GX 340+0. The directions in which inferred $\dot{M}$ increases are indicated by arrows.




Fig. 9.— *Top*: evolution of the 3 – 150 keV flux during the 2000 (*R5*) evolution events. Red/blue points A, B, and C mark moments at MJD = 51773.2/51774.7, 51773.6/51774.1 and 51772.2/51774.6 related to different phases of Z-state evolution. *Bottom*: PDSs for 15–30 keV band (*left* column) are plotted along with the $E * F(E)$ diagram (*right* column) for A, B and C points of X-ray light curve. $E * F(E)$ diagrams (panels A2, B2, C2) are related to the corresponding power spectra (panels A1, B1, C1). The data are shown by black points and the spectral model components are displayed by red, green, blue and dashed purple lines for *Comptb1*, *Comptb2*, *Blackbody* and *Gaussian* respectively.




Fig. 10.— *Left panel:* The $\nu \times power$ diagram of GX 340+0 in 0.01 – 150 Hz range, observed on August 18, 2000 (50016-01-02-16, MJD=51774.47, point D_1 on Fig. 9). The PDS accumulation time is 3876 sec. Blue solid line shows a smooth power spectrum. Broad QPO is centered at $\nu_{FBO} \sim 6$ Hz and very low frequency noise component (VLFN) are present during flaring branch (see details in the text). *Right panel:* $E * F(E)$ diagram is related to the corresponding power spectrum (point D_1). The data are shown by black points and the spectral model components are displayed by *red*, *green*, *blue* and *purple* dashed lines for *Comptb1*, *Comptb2*, *Blackbody* and *Gaussian* respectively. Yellow shaded area demonstrates relatively low contribution of *Comptb1* component at point D_1 .




Fig. 11.— A suggested geometry of GX 340+0 during state evolution. As GX 340+0 moves from HB, through NB, to FB. CC puffs up, which affects a visibility of the central area (NS). In the upper panel (FB) we schematically indicate a probable inclination of GX 340+0 as inferred from various observations (see text). In particular, the innermost part of the source is hidden by CC which causes dips in intensity (“dipping FB”). The successive decrease of screening effect during next stage (NB) is schematically shown in the middle panel. CC shrinking provides visibility of central region (NS). Finally, the lower (HB) panel demonstrates the minimal screening effect of central area, providing a proper contribution of NS emission.




Fig. 12.— L_{39} measured in $N_{COMPTB} \times D_{10}^2$ units versus electron temperature T_e (in keV) obtained for Z source GX 340+0 (*pink*), we present a variable N_{com1} normalization for this source; *atoll* sources GX 3+1 (*red*, taken from ST12) and 4U 1728-34 (*blue*, taken from ST11) for *RXTE* data. Here D_{10} is the distance to the source (see Table 5).